

### INITIAL RICE PROTEIN REFERENCES TO BE COMPLETED.

AACC (The American Association of Cereal Chemists),2000. Approved methods of American Association of Cereal Chemists. The Am. Assoc. Cereal Chem.Inc., St. Paul. Minnesota.

AOAC, 1990, Official Methods of Analysis, Association of Official Analytical Chemists (AOAC), Washington DC., 1 - 50.

Abdelrahman, S.M., H.B. Elmaki, W.H. Idris, A.B.Hassan, E.E. Babiker and A.H. El-Tinay, 2007. Antinutritional factor content and hydrochloric acid extractability of minerals in pearl millet cultivars as affected by germination. *Int. J. Food Sci. Nutr.*, 58: 6-17.

Aberle, T., Burchard, W., Vorweg, W. & Radosta, S. (1994). Conformational contributions of amylose and amylopectin to the structural properties of starches from various sources. *Starch/Stärke*, 46, 329–333.

Adam, J.M. (2006). Some quality factors of raw and processed rice. A paper presented at processing packing and marketing workshop W.D.C. Abakliki pg. 3-4.

Ahsan N, Komatsu S. Comparative analyses of the proteomes of leaves and flowers at various stages of development reveal organ specific functional differentiation of proteins in soybean. *Proteomics*. 2009;9(21):4889–4907. doi: 10.1002/pmic.200900308.

Alaka I.C. (2005). Upgrading the quality of local rice of for local and international market. A paper presented at 29th annual conference of Nigerian institute of food science and technology W.D.C. Abakaliki. Pg. 8.

Albritton DL, Derwent RG, Isaksen ISA, Lal M, Wuebbles DJ. In: *Climate change 1994 - radiative forcing of climate change and an evaluation of the IPCC IS92 Emission Scenarios*. Houghton JT, Meiro Filho LG, Bruce JP, Lee H, Callander BA, Haites EF, Harris N, Maskell K, editor. Cambridge University Press, Cambridge; 1994. Trace gas radiative forcing indices; pp. 205–246.

Ali M.A.M., Tinay A.H.E., Abdalla A.H. Effect of fermentation on the in vitro protein digestibility of pearl millet. *Food Chem*. 2003;80:51–54.

Ali, S.Z. Bhattacharya, K.R (1972) hydration and amylose solubility behavior of parboiled rice lebensm-wiss. *Technology* pg. 5, 207-212.

Ali, S.Z Bhattacharya, K.R (1980) changes of sugar and amino acids during parboiling of rice S. *Food biochem*, pg 4, 164-179.

Alladi S., R. Gilbert, and K. R. Sha nmugasundaram 1989. Lipids, lipoproteins id lipolytic activity in plasma with dietary protein changes. 40:653-661.

Almond R.J., Flanagan B.F., Antonopoulos A., Haslam S.M., Dell A., Kimber I., Dearman R.J. Differential immunogenicity and allergenicity of native and recombinant human lactoferrins: Role of glycosylation. *Eur. J. Immunol.* 2012;43:170–181.

Altintas MM, Rossetti MA, Nayer B, Puig A, Zagallo P, Ortega LM, Johnson KB, McNamara G, Reiser J, Mendez AJ, Nayer A. Apoptosis, mastocytosis, and diminished adipocytokine gene expression accompany reduced epididymal fat mass in long-standing diet-induced obese mice. *Lipids Health Dis.* 2011;10:198. doi: 10.1186/1476-511X-10-198.

Ananiadou S, Pyysalo S, Tsujii J, Kell DB. Event extraction for systems biology by text mining the literature. *Trends Biotechnol.* 2010;28:381–390. doi: 10.1016/j.tibtech.2010.04.005.

Anderson, A.K. and H.S. Guraya, 2001. Extractability of protein in physically processed rice bran. *J. Am. Oil Chem. Soc.*, 78: 969-972.

Anderson, J. W., and A. E. Siesel. 1990. Hypocholesterolemia effects of oat products, in *New Developments in Dietary Fiber: Physiological, Physicochemical, and Analytical* Furda, I., Brine, C.J., Eds., Plenum Press, New York, 17-36.

Anonymous. 1980. Official methods of analysis. (13th Ed.) Association of official analytical chemists. Arlington, USA. Bahnansey, Y., K. Khan and R. Harrold, 1986. Fortification of spaghetti with edible legumes. I. Physico-chemical, antinutritional, amino acid and mineral composition. *Cereal Chem.*, 63: 210-215.

Arai T, Takaya T, Ito Y, Hayakawa K, Toshima S, Shibuya C, Nomura M, Yoshimi N, Shibayama M, Yasuda Y. Bronchial asthma induced by rice. *Intern Med.* 1998;37:98–101. doi: 10.2169/internalmedicine.37.98. Asero R, Amato S, Alfieri B, Folloni S, Mistrello G. Rice: another potential cause of food allergy in patients sensitized to lipid transfer protein. *Int Arch Allergy Immunol.* 2007;143:69–74. doi: 10.1159/000098226.

Arai T, Takaya T, Ito Y, Hayakawa K, Toshima S, Shibuya C, et al. Bronchial asthma induced by rice. *Intern Med.* 1998;37:98–101.

Aranda B, Achuthan P, Alam-Faruque Y, Armean I, Bridge A, Derow C, Feuermann M, Ghanbarian AT, Kerrien S, Khadake J. et al. The IntAct molecular interaction database in 2010. *Nucleic Acids Res.* 2009;38:D525–531.

Arroyo V. Human serum albumin: Not just a plasma volume expander. *Hepatology.* 2009;50:355–357. Streatfield S.J., Howard J.A. Plant-based vaccines. *Int. J. Parasitol.* 2003;33:479–493.

Asadullah K., Sterry W., Volk H.D. Interleukin-10 therapy--review of a new approach. *Pharmacol Rev.* 2003;55:241–269.

Asano, H., Hirano, F., Isobe, K. & Sakurai, H. (2000). Effect of harvest time on the protein composition (glutelin, prolamin, albumin) and amylose content in paddy rice cultivated by Aigamo duck farming system. *Japanese Journal of Crop Science*, 69, 320–323.

Asero R, Amato S, Alfieri B, Folloni S, Mistrello G. Rice: another potential cause of food allergy in patients sensitized to lipid transfer protein. *Int Arch Allergy Immunol.* 2007;143:69–74.

Asero R, Mistrello G, Roncarolo D, Amato S, Caldironi G, Barocci F, et al. Immunological cross-reactivity between lipid transfer proteins from botanically unrelated plant-derived foods: a clinical study. *Allergy.* 2002;57:900–906.

Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, et al. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet.* 2000;25:25–29. doi: 10.1038/75556.

Assenov Y, Ramirez F, Schelhorn SE, Lengauer T, Albrecht M. Computing topological parameters of biological networks. *Bioinformatics.* 2008;24:282–284. doi: 10.1093/bioinformatics/btm554.

Association of Official Agricultural Chemists. *AOAC Official Methods of Analysis.* 16th ed. Chapter 4.

Association of Official Agricultural Chemists; Washington, DC, USA: 1995. p. 17.

Ausubel FM, Brent R, Kingston RE, Moore DD, Seidman JG, Smith JA, Struhl K. Phenol/SDS method for plant RNA preparation. John Wiley & Sons Inc; 1995.

Ayano. 1989. Effect of rice bran hemicellulose on the cholesterol metabolism in rats. *Nippon Eiyoshokuryo Gakkaishi.* 42:55-61.

Ayano Y., F. Ohta, Y. Watanabe, and K. Mita. 1980. Dietary fiber fractions in defatted rice bran and their hypocholesterolemia effect in cholesterol-fed rats.

Azhakanandam, K., Power, J.B., Lowe, K.C., et al. (2000). Qualitative assessment of aromatic Indica rice *Oryza sativa* L. Proteins, lipids and starch in grain from somatic embryo- and seed-derived plants. *Journal of Plant Physiology,* 156, 783–789.

B. O. Eggum, B. O. Juliano, and C. C. Mangingat, 1982, Protein and energy utilization of rice milling fractions, *Journal of Human Nutrition,* 31, 371 – 376.

Bailey C.J., Turner R.C. Metformin. *New Eng. J. Med.* 1996;334:574–579.

Baler M, Dietz KJ. The plant 2-Cys peroxiredoxin BAS1 is a nuclear-encoded chloroplast protein: its expressional regulation, phylogenetic origin, and implications for its specific physiological function in plants. *Plant J.* 1997;12(1):179–190. doi: 10.1046/j.1365-3113.1997.12010179.x.

Banks, W., Greenwood, C.T. & Muir, D.D. (1974). The characterisation of starch and its components. *Starch/Starke,* 26, 73–103.

Bao, J.S., He, P., Xia, Y.W., Chen, Y. & Zhu, L.H. (1999). Starch RVA profile parameters of rice are mainly controlled by Wx gene. *Chinese Science Bulletin,* 44, 2047–2051.

- Bao, J.S., Sun, M. & Corke, H. (2002). Analysis of the genetic behavior of some starch properties in indica rice (*Oryza sativa* L.): thermal properties, gel texture, swelling volume. *Theoretical and Applied Genetics*, 104, 408–413.
- Barber, S. (1972) milled rice and changes during ageing in the chemistry and tech 1st eds (D.F. Houston Ed) AM Associate. *Cereal chem.* St. Paul, MN U. S. A. pg. 215-263.
- Barber, S. (1972). Milled rice and changes during aging. In: *Rice: Chemistry and Technology* (edited by D.F. Houston). Pp. 215–263. St. Paul, Minnesota: American Association of Cereal Chemists.
- Bardor M., Faveeuw C., Fitchette A.C., Gilbert D., Galas L., Trottein F., Faye L., Lerouge P. Immunoreactivity in mammals of two typical plant glyco-epitopes, core alpha(1,3)-fucose and core xylose. *Glycobiology*. 2003;13:427–434.
- Barnett C.R., Flatt P.R., Ioannides C. Modulation of the rat hepatic cytochrome P450 composition by long-term streptozotocin-induced insulin-dependent diabetes. *J. Biochem. Toxicol.* 1994;2:63–69.
- Barta A., Sommergruber K., Thompson D., Hartmuth K., Matzke M.A., Matzke A.J.M. The expression of a napoline synthase human growth hormone chimeric gene in transformed tobacco and sunflower callus tissue. *Plant Mol. Biol.* 1986;6:347–357.
- Basak, S., Tyagi, R.S. & Srivastava, K.N. (2002). Biochemical characterization of aromatic and non-aromatic rice cultivars. *Journal of Food Science and Technology – Mysore*, 39, 55–58.
- Bates, F.L., French, D. & Rundle, R.G. (1943). Amylose and amylopectin content of starches determined by their complex formation. *Journal of the American Chemical Society*, 65, 142–148.
- Batey, I.L. & Curtin, B.M. (1996). Measurement of amylose/amylopectin ratio by high-performance liquid chromatography. *Starch/Stärke*, 48, 338–344.
- Bean, M.M. and K.D. Nishita. 1985. Rice flours for baking. In: *Rice Chemistry and Technology*. (Ed.): B.O. Juliano. AACC International, St. Paul, MN, pp. 539-556.
- Bechtel, D.B. & Pomeranz, Y. (1977). Ultrastructure of the mature ungerminated rice (*Oryza sativa*) caryopsis. The caryopsis coat and aleurone cells. *American Journal of Botany*, 64, 966–973.
- Bechtel, D.B. & Pomeranz, Y. (1978). Ultrastructure of the mature ungerminated rice (*Oryza sativa*) caryopsis. The starchy endosperm. *American Journal of Botany*, 65, 684–688.
- Begum, M. & Bhattacharya, R. (2000). Fat contents in raw and cooked Assamrice varieties. *Journal of Food Science and Technology – Mysore*, 37, 58–60.
- Bello-Perez, L.A., Roger, P., Baud, B. & Colonna, P. (1998). Macromolecular features of starches determined by aqueous high-performance size exclusion chromatography. *Journal of Cereal Science*, 27, 267–278.

- Benedito-de, B.C. and S. Barber, 1978. Toxic constituents of rice bran: Haemagglutinating activity of raw and heat treated bran. *Rev. Agroquim. Technol. Aliment.*, 18: 89-94.
- Bent AF. Arabidopsis in planta transformation. Uses, mechanisms, and prospects for transformation of other species. *Plant Physiol.* 2000;124:1540–1547. doi: 10.1104/pp.124.4.1540.
- Bera, M.B. and R.K. Mukherjee, 1988. Nutritional evaluation of rice bran protein concentrate. *Ind. J. Nutr. Diet.*, 25: 50-54.
- Bera, M.B. and R.K. Mukherjee, 1989. Solubility, emulsifying and foaming properties of rice bran protein concentrates. *J. Food Sci.*, 54: 142-145.
- Berglund AC, Sjolund E, Ostlund G, Sonnhammer EL. InParanoid 6: eukaryotic ortholog clusters with inparalogs. *Nucleic Acids Res.* 2008;36:D263–266.
- Bergman, C.J., Delgado, J.T., Bryant, R., Grimm, C., Cadwallader, K.R. & Webb, B.D. (2000). Rapid gas chromatographic technique for quantifying 2-acetyl-1-pyrroline and hexanal in rice (*Oryza sativa* L.). *Cereal Chemistry*, 77, 454–458.
- Berry J.D., Gaudet R.G. Antibodies in infectious diseases: Polyclonals, monoclonals and niche biotechnology. *New Biotechnol.* 2011;28:489–501.
- Bevan M.W., Flavell R.B., Chilton M.D. A chimaeric antibiotic resistance gene as a selectable marker for plant cell transformation. 1983. *Biotechnology.* 1992;24:367–370.
- Bhattacharjee, P., Singhal, R.S. & Kulkarni, P.R. (2002). Basmati rice: a review. *International Journal of Food Science and Technology*, 37, 1–12.
- Bhattacharya, K.R Sowbhagy A.C.M (1971) water uptake by rice drying cooking cereal science today vol. 6 pg. 420-424.
- Bhattacharya, K.R. Sowbhagy A.C. and Indudhara Swamy. Y.M. (1978) importance of insoluble amylase determinant of rice quality journal science food Agri vol 29 pg. 359-364.
- Biliaderis, C.G. & Tonogai, J.R. (1991). Influence of lipids on the thermal and mechanical properties of concentrated starch gels. *Journal of Agricultural and Food Chemistry*, 39, 833–840.
- Binns D, Dimmer E, Huntley R, Barrell D, O'Donovan C, Apweiler R. QuickGO: a web-based tool for Gene Ontology searching. *Bioinformatics.* 2009;25:3045–3046. doi: 10.1093/bioinformatics/btp536.
- Biolo G, Tipton KD, Klein S, Wolfe RR: An abundant supply of amino acids enhances the metabolic effect of exercise on muscle protein. *Am J Physiol* 1997, 273:E122-E129.
- Birmingham NP, Parvataneni S, Hassan HM, Harkema J, Samineni S, Navuluri L, Kelly CJ, Gangur V. An adjuvant-free mouse model of tree nut allergy using hazelnut as a model tree nut. *Int Arch Allergy Immunol.* 2007;144:203–210. doi: 10.1159/000103993.

- Biswas, S.K. and B.O. Juliano, 1988. Laboratory parboiling procedures and properties of parboiled rice from varieties differing in starch properties. *Cereal Chem.*, 65: 417-423.
- Blakeney, A. (1984). Rice grain quality. In: *Rice Growing in New South Wales* (edited by A. Currey). Pp. 1–5. Yanco, Australia: Department of Agriculture New South Wales and the Rice Research Committee.
- Blank C.A., Brantly M. Clinical features and molecular characteristics of alpha 1-antitrypsin deficiency. *Ann. Allergy*. 1994;72:105–120. ; quiz 120–102.
- Blanshard, J.M.V. (1987). Starch granule structure and function: a physicochemical approach. In: *Starch: Property and Potential* (edited by T. Galliard). Pp. 16–54. Chichester: John Wiley.
- Blomstrand E: A role for branched-chain amino acids in reducing central fatigue. *J Nutr* 2006, 136:544S-547S.
- Blum, H., H. Beier and H.J. Gross. 1987. Improved silver staining of plant proteins, RNA and DNA in polyacrylamide gels. *Electrophoresis*, 8: 93-99.
- Boivin E.B., Lepage E., Matton D.P., De Crescenzo G., Jolicoeur M. Transient expression of antibodies in suspension plant cell suspension cultures is enhanced when co-transformed with the tomato bushy stunt virus p19 viral suppressor of gene silencing. *Biotechnol. Prog.* 2010;26:1534–1543.
- Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem.* 1976;72:248–254. doi: 10.1016/0003-2697(76)90527-3.
- Blom N, Sicheritz-Ponten T, Gupta R, Gammeltoft S, Brunak S. Prediction of post-translational glycosylation and phosphorylation of proteins from the amino acid sequence. *Proteomics*. 2004;4(6):1633–1649. doi: 10.1002/pmic.200300771.
- Bortesi L., Rossato M., Schuster F., Raven N., Stadlmann J., Avesani L., Falorni A., Bazzoni F., Bock R., Schillberg S., et al. Viral and murine interleukin-10 are correctly processed and retain their biological activity when produced in tobacco. *BMC Biotechnol.* 2009;9:22.
- Brand-Miller J., Hayne S., Petocz P., Colagiuri S. Low-glycemic index diets in the management of diabetes: A meta-analysis of randomized controlled trials. *Diabetes Care.* 2003;26:2261–2267.
- Brandao MM, Dantas LL, Silva-Filho MC. AtPIN: Arabidopsis thaliana protein interaction network. *BMC Bioinformatics.* 2009;10:454. doi: 10.1186/1471-2105-10-454.
- Brandsch C, Schmidt T, Behn D, Weibe K, Mueller AS, Stangl G: Glutathione deficiency down-regulates hepatic lipogenesis in rats. *Lipids Health Dis* 2010, 9:50.
- Bray GA. Obesity: the disease. *J Med Chem.* 2006;49:4001–4007. doi: 10.1021/jm0680124.

Brears T, Liu C, Knight TJ, Coruzzi GM. Ectopic overexpression of asparagine synthetase in transgenic tobacco. *Plant Physiol.* 1993;103:1285–1290.

Brosnan JT, Brosnan ME: Branched-chain amino acids: enzyme and substrate regulation. *J Nutr* 2006, 136:207S-211S.

Browman DT, Hoegg MB, Robbins SM. The SPFH domain-containing proteins: more than lipid raft markers. *Trends Cell Biol.* 2007;17:394–402. doi: 10.1016/j.tcb.2007.06.005.

Brown KR, Jurisica I. Unequal evolutionary conservation of human protein interactions in interologous networks. *Genome Biol.* 2007;8:R95. doi: 10.1186/gb-2007-8-5-r95.

Brown KR, Jurisica I. Online predicted human interaction database. *Bioinformatics.* 2005;21:2076–2082. doi: 10.1093/bioinformatics/bti273.

Buleon, A., Colonna, P., Planchot, V. & Ball, S. (1998). Starch granules-structure and biosynthesis. *International Journal of Biological Macromolecules*, 23, 85–112.

Bunney TD, van Walraven HS, de Boer AH. 14-3-3 protein is a regulator of the mitochondrial and chloroplast ATP synthase. *Proc Natl Acad Sci USA.* 2001;98(7):4249–4254. doi: 10.1073/pnas.061437498.

Butland G, Peregrin-Alvarez JM, Li J, Yang W, Yang X, Canadien V, Starostine A, Richards D, Beattie B, Krogan N. et al. Interaction network containing conserved and essential protein complexes in *Escherichia coli*. *Nature.* 2005;433:531–537. doi: 10.1038/nature03239.

Butterworth, R.F., L. Spahr, S. Fontaine, and P. Layragues, Manganese Toxicity, Dopaminergic Dysfunction, and Hepatic Encephalopathy, *Metab. Brain Dis.* 10:259–267 (1995).

Byvatov E, Schneider G. Support vector machine applications in bioinformatics. *Appl Bioinformatics.* 2003;2:67–77.

C. K. Probart, P.J. Bird, and K. A. Parker, 1993, Diet and Athletic Performance. *Medicine and Clinical Journal of North America*, 5, 77- 757.

C. O. Edeogu, F. C. Ezeonu, A. N. C. Okaka, C. E. Ekuma, and S. O. Elom, 2007, Proximate composition of staple food crops in Ebonyi State (South Eastern Nigeria), *International Journal of Biotechnology and Biochemistry*, 3(1), 1-8.

C. Eromosele and C. O. Eromosele, 1994, Studies on the chemical composition and physico-chemical properties of seeds of some wild plants, *Plant Foods for Human Nutrition*, 46, 361-365.

C. S. James, 1995, Mineral elements in rice, *Analytical Chemistry of Food*, Blackie Academic and Professional Publishers, New York, 126 – 128.

Cacas JL. Devil inside: does plant programmed cell death involve the endomembrane system? *Plant Cell Environ.* 2010;33:1453–1473.

- Cagampang, G., L. Cruz, S. Espiritu, R. Santiago and B.O. Juliano. 1966. Studies on the extraction and composition of rice proteins. *Cereal Chem.*, 43: 145-155.
- Cagampang, G.B., L.J. Cruz, S.G. Espiritu, R.G. Santiago, and B.O. Juliano, Studies on the Extraction and Composition of Rice Proteins, *Cereal Chem.* 43:145–155 (1966).
- Calvert, C., K. Parker, Z. J. Parker, R. N. Sayre, and R. M. Saunders. 1985. Rice bran in swine rations. *Calif. Agric.* May-June, 19.
- Camou JP, Sebranek, JG. "Effect of non-meat proteins on gelation properties of porcine muscle proteins." *Journal of Muscle Foods.* 1991; 2: 149-163. Food and Nutrition Press Inc., Trumbull, CT.
- Campbell B, Kreider RB, Ziegenfuss T, La Bounty P, Roberts M, Burke D, Landis J, Lopez H, Antonio J: International society of sports nutrition position stand: protein and exercise. *J Int Soc Sports Nutr* 2007, 4:8.
- Capobianco F, Butteroni C, Barletta B, Corinti S, Afferni C, Tinghino R, Boirivant M, Di Felice G. Oral sensitization with shrimp tropomyosin induces in mice allergen-specific IgE, T cell response and systemic anaphylactic reactions. *Int Immunol.* 2008;20:1077–1086. doi: 10.1093/intimm/dxn065.
- Caraco Y., Muszkat M., Wood A.J. Phenytoin metabolic ratio: A putative marker of CYP2C9 activity in vivo. *Pharmacogenetics.* 2001;11:587–596.
- Carter C, Pan S, Zouhar J, Avila EL, Girke T, Raikhel NV. The vegetative vacuole proteome of *Arabidopsis thaliana* reveals predicted and unexpected proteins. *Plant Cell.* 2004;16:3285–3303. doi: 10.1105/tpc.104.027078.
- Carter CJ, Bednarek SY, Raikhel NV. Membrane trafficking in plants: new discoveries and approaches. *Curr Opin Plant Biol.* 2004;7:701–707. doi: 10.1016/j.pbi.2004.09.016.
- Cavalier D.M., Lerouxel O., Neumetzler L., Yamauchi K., Reinecke A., Freshour G., Zobotina O.A., Hahn M.G., Burgert I., Pauly M., et al. Disrupting two *Arabidopsis thaliana* xylosyltransferase genes results in plants deficient in xyloglucan, a major primary cell wall component. *Plant. Cell.* 2008;20:1519–1537.
- Ceol A, Chatr Aryamontri A, Licata L, Peluso D, Briganti L, Perfetto L, Castagnoli L, Cesareni G. MINT, the molecular interaction database: 2009 update. *Nucleic Acids Res.* 2010;38:D532–539. doi: 10.1093/nar/gkp983.
- CeraLyte®/CeraSport® Assures Safety of Its Rice-Based Products. Only California rice used in its superior oral rehydration solution and hydration drinks (news/article/1211707/CeraLyte-ceraSport-Assures-Safety-of-its-Rice-Based-products-Only-California-rice).
- Champagne, E.T., Bett, K.L., Vinyard, B.T., et al. (1999). Correlation between cooked rice texture and Rapid Visco Analyses measurements. *Cereal Chemistry*, 76, 764–771.
- Chandi, G.K. and D.S. Sogi, 2007. Functional properties of rice bran protein concentrates *J. Food Eng.* 79:



592-597.

Chang, S.M. & Chang, Y.C. (1988). Retrogradation of rice starches with various amylose contents. *Proceedings of the National Science Council*, 12, 247–251.

Chang, S.M. & Liu, L.C. (1991). Retrogradation of rice starches studied by differential scanning calorimetry and influence of sugars, NaCl and lipids. *Journal of Food Science*, 56, 564–567.

Chatakanonda, P., Varavinit, S. & Chinachoti, P. (2000). Effect of crosslinking on thermal and microscopic transitions of rice starch. *Food Science and Technology*, 33, 276–284.

Chatr-aryamontri A, Ceol A, Palazzi LM, Nardelli G, Schneider MV, Castagnoli L, Cesareni G. MINT: the Molecular INTERaction database. *Nucleic Acids Res.* 2007;35:D572–574. doi: 10.1093/nar/gkl950.

Chavan, J. and S. Duggal. 1978. Studies on the essential amino acid composition, protein fractions and biological value (BV) of some new varieties of rice. *J. Sci. Food Agri.*, 29: 225-229.

Chen F, Yuan Y, Li Q, He Z. Proteomic analysis of rice plasma membrane reveals proteins involved in early defense response to bacterial blight. *Proteomics.* 2007;7:1529–1539. doi: 10.1002/pmic.200500765.

Chen JY, Youn E, Mooney SD. Connecting protein interaction data, mutations, and disease using bioinformatics. *Methods Mol Biol.* 2009;541:449–461. doi: 10.1007/978-1-59745-243-4\_19.

Chen T.L., Lin Y.L., Lee Y.L., Yang N.S., Chan M.T. Expression of bioactive human interferon-gamma in transgenic rice cell suspension cultures. *Transgenic Res.* 2004;13:499–510.

Chen X.J., Guo Z.J. Tobacco OPBP1 enhances salt tolerance and disease resistance of transgenic rice. *Int. J. Mol. Sci.* 2008;9:2601–2613.

Chen, Y., Y. Chen, C. Wu, C. Yu and H. Liao. 2010. Prolamin, a rice protein, augments anti-leukaemia immune response. *J. Cereal Sci.*, 51: 189-197.

Chen Z.Y., Jiao R., Ma K.Y. Cholesterol-lowering nutraceuticals and functional foods. *J. Agric. Food Chem.* 2008;56:8761–8773.

Cheng HC, Cheng PT, Peng P, Lyu PC, Sun YJ. Lipid binding in rice nonspecific lipid transfer protein-1 complexes from *Oryza sativa*. *Protein Sci.* 2004;13:2304–2315.

Cheung MY, Zeng NY, Tong SW, Li FWY, Zhao KJ, Zhang Q, Sun SSM, Lam HM. Expression of a RING-HC protein from rice improves resistance to *Pseudomonas syringae* pv. tomato DC3000 in transgenic *Arabidopsis thaliana*. *J Exp Bot.* 2007;58:4147–4159. doi: 10.1093/jxb/erm272.

Cheung MY, Zeng NY, Tong SW, Li FWY, Xue Y, Zhao KJ, Wang C, Zhang Q, Fu Y, Sun Z. et al. Constitutive expression of a rice GTPase-activating protein induces defense responses. *New Phytol.* 2008;179:530–545. doi: 10.1111/j.1469-8137.2008.02473.x.

Chiang, P.Y. & Yeh, A.I. (2002). Effect of soaking on wetmilling of rice. *Journal of Cereal Science*, 35, 85–94. Composition and functional properties of rice Z. Zhou et al. 863 \_ 2002 Blackwell Science Ltd  
*International Journal of Food Science and Technology* 2002, 37, 849–868.

Chien, J.T., Lien, Y.Y. & Shoemaker, C.E. (1999). Effect of polarity of complexing agents on thermal and rheological properties of rice starch gels. *Cereal Chemistry*, 76, 837–842.

Chinnaswamy, R. & Bhattacharya, K.R. (1986). Characteristics of gel chromatographic fractions of starch in relation to rice and expanded rice-product qualities. *Starch/Starke*, 38, 51–57.

Chitteti BR, Peng Z. Proteome and phosphoproteome differential expression under salinity stress in rice (*Oryza sativa* L.) roots. *J Proteome Res.* 2007;6(5):1718–1727. doi: 10.1021/pr060678z.

Chitteti BR, Peng Z. Proteome and phosphoproteome dynamic change during cell dedifferentiation in *Arabidopsis*. *Proteomics.* 2007;7(9):1473–1500. doi: 10.1002/pmic.200600871.

Cho J.S., Hong S.M., Joo S.Y., Yoo J.S., Kim D.I. Cryopreservation of transgenic rice suspension cells producing recombinant hCTLA4Ig. *Appl. Microbiol. Biotechnol.* 2007;73:1470–1476.

Choi MH, Lee IK, Kim GW, Kim BU, Han YH, Yu DY, Park HS, Kim KY, Lee JS, Choi C, Bae YS, Lee BI, Rhee SG, Kang SW. Regulation of PDGF signalling and vascular remodelling by peroxiredoxin II. *Nature.* 2005;435(7040):347–353. doi: 10.1038/nature03587.

Choudhury, N.H. & Juliano, B.O. (1980a). Lipids in developing and mature rice grain. *Phytochemistry*, 19, 1063–1069.

Choudhury, N.H. & Juliano, B.O. (1980b). Effect of amylose content on the lipids of mature rice grain. *Phytochemistry*, 19, 1385–1389.

Chrastil, J. 1992. Correlations between the physicochemical and functional properties of rice. *J. Agri. Food Chem.*, 40:1683-1686.

Chrastil J. (1994) effected of storage on the physiochemical properties and quality factors of rice. In rice scenes and technology (W.E. Maradell and J.I. Wads Worth, eds), Marceldeker, new tark pg. 49-81.

Chrastil J. (1990) chemical and physiochemical changes of rice during storage at different temperature *Journal of cereal vol ii* pg. 75-85.

Chrastil (1990) protein-starch interaction in rice grains influence of storage on oryzenin and starch. *Journal. Agric food chem Vol 38* pg. 184-189. Chrastil, J. (1990). Chemical and physicochemical changes of rice during storage at different temperatures. *Journal of Cereal Science*, 11, 71–85.

Chun WS, Lee KS, Hong CH, Lee SY. A study on the cross-allergenicity between buckwheat and rice flour using IgE-immunoblot inhibition and ELISA-inhibition test. *Pediatr Allergy Respir Dis.* 2000;10:161–170.

Chung, H.J., Lee, E.J. & Lim, S.T. (2002). Comparison in glass transition and enthalpy relaxation between native and gelatinized rice starches. *Carbohydrate Polymers*, 48, 287–298.

Chung E, Oh SK, Park JM, Choi D. Expression and promoter analyses of pepper CaCDPK4 (*Capsicum annuum* calcium dependent protein kinase 4) during plant defense response to incompatible pathogen. *Plant Pathol J*. 2007;23:76–89. doi: 10.5423/PPJ.2007.23.2.076.

Chung, O. and Y. Pomeranz. 1985. Amino acids in cereal proteins and protein fractions. Digestibility and amino acid availability in cereals and oilseeds, pp. 65-107.

Claros MG, Heijne G. TopPred II: an improved software for membrane protein structure predictions. *Comp Appl Biosci*. 1994;10:685–686.

Connor, M. A., R. M. Saunders, G. O. Kohler. 1976. Rice bran protein concentrates obtained by wet alkaline extraction. *Cereal Chem*. 53:488-496.

Cornell, H.J., McGrane, S.J. & Rix, C.J. (1999). A novel and rapid method for the partial fractionation of starch using 1-butanol in the presence of thiocyanate. *Starch/Starke*, 51, 335–341.

Cui J, Li P, Li G, Xu F, Zhao C, Li Y, Yang Z, Wang G, Yu Q, Shi T. AtPID: Arabidopsis thaliana protein interactome database--an integrative platform for plant systems biology. *Nucleic Acids Res*. 2008;36:D999–1008.

Daniel (1998): effect of long grain rice storage history and end quality journal food sciences 63 vol (5) pg. 832.

Das UN: Essential fatty acids and their metabolites could function as endogenous HMG-CoA reductase and ACE enzyme inhibitors, antiarrhythmic, anti-hypertensive, anti-atherosclerotic, anti-inflammatory, cytoprotective, and cardioprotective molecules. *Lipids Health Dis* 2008, 7:37.

Davis JM: Carbohydrates, branched-chain amino acids, and endurance: the central fatigue hypothesis. *Int J Sport Nutr* 1995, 5(Suppl):S29-S38.

Dawson P.A., Rudel L.L. Intestinal cholesterol absorption. *Curr. Opin. Lipidol*. 1999;10:315–320.

Dearman RJ, Kimber I. Animal models of protein allergenicity: potential benefits, pitfalls and challenges. *Clin Exp Allergy*. 2009;39:458–468. doi: 10.1111/j.1365-2222.2008.03194.x.

Dearman RJ, Kimber I. A mouse model for food allergy using intraperitoneal sensitization. *Methods*. 2007;41:91–98. doi: 10.1016/j.ymeth.2006.07.008.

Dearman RJ, Stone S, Caddick HT, Basketter DA, Kimber I. Evaluation of protein allergenic potential in mice: dose-response analyses. *Clin Exp Allergy*. 2003;33:1586–1594. doi: 10.1046/j.1365-2222.2003.01793.x.

De Bodt S, Proost S, Vandepoele K, Rouze P, Van de Peer Y. Predicting protein-protein interactions in *Arabidopsis thaliana* through integration of orthology, gene ontology and co-expression. *BMC Genomics*. 2009;10:288. doi: 10.1186/1471-2164-10-288.

De Castro E, Sigrist CJA, Gattiker A, Bulliard V, Langendijk-Genevaux PS, Gasteiger E, Bairoch A, Hulo N. ScanProsite: detection of PROSITE signature matches and ProRule-associated functional and structural residues in proteins. *Nucl Acids Res*. 2006;34:W362–W365. doi: 10.1093/nar/gkl124.

De Neve M., De Buck S., de Wilde C., van Houdt H., Strobbe I., Jacobs A., van Montagu M., Depicker A. Gene silencing results in instability of antibody production in transgenic plants. *Mol. Gen. Genet*. 1999;260:582–592.

De S, Babu MM. Genomic neighbourhood and the regulation of gene expression. *Curr Opin Cell Biol*. 2010;22:326–333. doi: 10.1016/j.ceb.2010.04.004.

Decker, L.A., 1977. *Worthington enzyme manual*. Worthington Biochemical Corp, Feedhold, NJ.

Deka, S.C., Sood, D.R. & Gupta, K.R. (2000). Nutritional evaluation of basmati rice (*Oryza sativa* L.) genotypes. *Journal of Food Science and Technology – Mysore*, 37, 272–276.

Deka S.C Sood D.R. and Gupta. S.K. (2000) effect of storage on fatty acid profile of basmati rice (*oryza sativa*) genotypes.

Delom F, Chevet E. Phosphoprotein analysis: from proteins to proteomes. *Proteome sci*. 2006;4:15. doi: 10.1186/1477-5956-4-15.

Del Rosario, A.R., Briones, V.P., Vidal, A.J. & Juliano, B.O. (1968). Composition and endosperm structure of developing and mature rice kernel. *Cereal Chemistry*, 45, 225–235.

Delwiche, S.R., Bean, M.M., Miller, R.E., Webb, B.D. & Williams, P.C. (1995). Apparent amylose content of milled rice by near-infrared reflectance spectrophotometry. *Cereal Chemistry*, 72, 182–187.

Deolankar, R.P. and K.S. Singh, 1979. Trypsin inhibitor, mineral availability and performance of broiler chickens fed on diets based on rice bran. *Anim. Feed Sci. Technol.*, 4: 133-141.

Diets with High or Low Protein Content and Glycemic Index for weight-Loss Maintenance. *New England Journal of Medicine*, 2010; 363:2102-13.

Di Lernia V, Albertini G, Bisighini G. Immunologic contact urticaria syndrome from raw rice. *Contact Dermatitis*. 1992;27:196. doi: 10.1111/j.1600-0536.1992.tb05260.x.

Dimopoulos J.S. Muller, H.G. (1972) effect of processing conditions on protein extraction and composition and on some other physiochemical characteristic of parboiled rice cereal chem. Vol 49 pg. 54-61.

DoctorsHealthPress.com Reports on Study: Vinegar That Helps Keep Diabetes Under Control (news article/1190001/DoctorsHealthPress.com Reports-on-Study-Vinegar-That-Helps-Keep-Diabetes Under Control (news/article/1190001/DoctorsHealthPress.com, Reports-on-Study-Vinegar-That-Helps-Keep-Diabetes-Under-Control)).

Dubois, M., K.A. Gilles, J.K. Hamilton, and F. Smith, Calorimetric Method for Determination of Sugar and Related Substances, *Anal. Chem.* 28:350–356 (1956).

E. N. Onyeike, T. Olungwe, and A. A. Uwakwe, 1995, Effect of heat treatment and defatting on the proximate composition of some Nigerian local soup thickeners, *Food Chemistry*, 53, 173-175.

E. O. Ibukun, 2008, Effect of prolonged parboiling duration on proximate composition of rice, *Journal of Scientific Research and Essay*, 3 (7), 323 – 325.

Easterling DR, Horton B, Jones PD, Peterson TC. Maximum and minimum temperature trends for the globe. *Science*. 1997;277:364–367. doi: 10.1126/science.277.5324.364.

Edenberg H.J. The genetics of alcohol metabolism: Role of alcohol dehydrogenase and aldehyde. *Alcohol. Res. Health*. 2007;30:5–13.

Ellis, R.P., Cochrane, M.P., Dale, M.F.B., et al. (1998). Starch production and industrial use. *Journal of the Science of Food and Agriculture*, 77, 289–311. Fan, J. & Marks, B.P. (1999). Effects of rough rice storage conditions on gelatinization and retrogradation properties of rice flours. *Cereal Chemistry*, 76, 894–897.

Enrique E, Ahrazem O, Bartra J, Latorre MD, Castelló JV, de Mateo JA, et al. Lipid transfer protein is involved in rhinoconjunctivitis and asthma produced by rice inhalation. *J Allergy Clin Immunol*. 2005;116:926–928.

Extracting Rice Protein with Different Proteases (J): *Cereal & Feed Industry*: 2002-02.

Fagerhol M.K., Laurell C.B. The polymorphism of “prealbumins” and alpha-1-antitrypsin in human sera. *Clin. Chim. Acta*. 1967;16:199–203.

Fan, J., Marks, B.P., Daniels, M.J. & Siebenmorgen, T.J. (1999). Effects of postharvest operations on the gelatinization and retrogradation properties of long-grain rice. *Transactions of the ASAE*, 42, 727–731.

Farran I., Sanchez-Serrano J.J., Medina J.F., Prieto J., Mingo-Castel A.M. Targeted expression of human serum albumin to potato tubers. *Transgenic Res*. 2002;11:337–346.

FDA Looks for Answers on Arsenic in Rice (news article 1172936/FDA-Looks-for-Answers-on-Arsenic-in-Rice).

F. E. Nwilene, S. O. Oikeh, T. A. Agunbiade, O. Oladimeji, O. Ajayi, M. Sie, G. B. Gregorio, A. Togota, and A. D. Taura, 2007, Increasing rice production in sub-Saharan Africa, challenges and opportunities, A production hand book, Africa.

Fell, J.M.E., A.P. Reynolds, N. Meadows, K. Khan, S.G. Long, G. Quaghebeur, W.J. Taylor, and J.P. Milla, Manganese Toxicity in Children Receiving Long-Term Parenteral Nutrition, *Lancet* 347:1218–1221 (1996).

Figueredo E, Quirce S, del Amo A, Cuesta J, Arrieta I, Lahoz C, et al. Beer-induced anaphylaxis: identification of allergens. *Allergy*. 1999;54:630–634.

Finckh U, Lingenfelter PA, Myerson D. Producing single-stranded DNA probes with the Taq DNA polymerase: a high yield protocol. *BioTechniques*. 1991;10:35–36. 38-39.

Fiocchi A, Bouygue GR, Restani P, Gaiaschi A, Terracciano L, Martelli A. Anaphylaxis to rice by inhalation. *J Allergy Clin Immunol*. 2003;111:193–195.

First Double Blind Study Proves Plant-based Rice Protein has Identical Benefits to Animal-based Whey Protein (News article/1349955/First-Double-Blind-Study-Proves-Plant-based-Rice-Protein-Has-Identical-Benefits).

Folch J, Lees M, Sloane-Stanley GH: A simple method for the isolation and purification of total lipids from animal tissues. *J Biol Chem* 1957, 226:497-509.

Food and Agriculture Organization / International Rice Research Institute, 2006, *FAO Food and Nutrition Series*, FAO Rome, 26 .

Food and Agriculture Organization / World Health Organization, 1998, *Obesity: Preventing and managing global epidemic*, WHO technical report, Geneva, Switzerland,11-12.

Foyer CH. Stromal protein phosphorylation in spinach (*Spinacia oleracea*) chloroplasts. *Biochem J*. 1985;231(1):97–103.

Franconi R., Massa S., Illiano E., Mullar A., Cirilli A., Accardi L., Di Bonito P., Giorgi C., Venuti A. Exploiting the plant secretory pathway to improve the anticancer activity of a plant-derived HPV16 E7 vaccine. *Int. J. Immunopathol. Pharmacol*. 2006;19:187–197.

Freeman J., Sparks C.A., West J., Shewry P.R., Jones H.D. Temporal and spatial control of transgene expression using a heat-inducible promoter in transgenic wheat. *Plant. Biotechnol. J*. 2011;9:788–796.

Friedman, M. and D.L. Brandon, 2001. Nutritional and health benefits of soy proteins. *J. Agric. Food Chem.*, 49: 1069-1086.

Fu TJ, Abbott UR, Hatzos C. Digestibility of food allergens and nonallergenic proteins in simulated gastric fluid and simulated intestinal fluid-a comparative study. *J Agric Food Chem*. 2002;50:7154–7160.

Fujino, Y. & Mano, Y. (1972). Classification of lipids and composition of fatty acids in brown rice. *Eiyo to Shokuryo*, 25, 472–474.

Fujita, S., Morita, T. & Fujiyama, G. (1993). The study of melting temperature and enthalpy of starch from rice, barley, wheat, foxtail-millet and proso-millet. *Starch/Starke*, 45, 436–441.

Fujiwara Y., Aiki Y., Yang L., Takaiwa F., Kosaka A., Tsuji N.M., Shiraki K., Sekikawa K. Extraction and purification of human interleukin-10 from transgenic rice seeds. *Protein Expr. Purif.* 2010;72:125–130.

Furukawa, S., T. Mizuma, Y. Kiyokawa, T. Masumura, K. Tanaka and Y. Wakai. 2003. Distribution of storage proteins in low-glutelin rice seed determined using a fluorescent antibody. *J. Biosci. Bioengin.*, 96: 467-473.

G. M. Wardlaw and M. Kessel, 2002, *Prospective in Nutrition*, 5th Edition, McGraw-Hill, Boston, 278.

Gabriel G, Limas MS, Moneo Ignacio, Fischer Stefan, Wittmann-Liebold Brigitte, Méndez Enrique. Purification and characterization of ten new rice NaCl-soluble proteins: identification of four protein-synthesis inhibitors and two immunoglobulin-binding proteins. *Planta*. 1990;181:1–9.

Garlick PJ: The role of leucine in the regulation of protein metabolism. *J Nutr* 2005, 135:1553S-1556S.

Garoosi G.A., Salter M.G., Caddick M.X., Tomsett A.B. Characterization of the ethanol-inducible alc gene expression system in tomato. *J. Exp. Bot.* 2005;56:1635–1642.

Geisler-Lee J, O'Toole N, Ammar R, Provart NJ, Millar AH, Geisler M. A predicted interactome for *Arabidopsis*. *Plant Physiol.* 2007;145:317–329. doi: 10.1104/pp.107.103465.

Gerhardt A. L., and N. B. Gallo. 1998. Full-fat rice bran and oat bran similarly reduce hypercholesterolemia in humans. *J. Nutr* 128:865-869.

Gibbons GF: Regulation of fatty acid and cholesterol synthesis: cooperation or competition? *Prog Lipid Res* 2003, 42:479-497.

Gilbert G, Lees A: Changes in the force development characteristics of muscle following repeated maximum force and power exercise. *Ergonomics* 2005, 48:1576-1584.

Giot L, Bader JS, Brouwer C, Chaudhuri A, Kuang B, Li Y, Hao YL, Ooi CE, Godwin B, Vitols E. et al. A protein interaction map of *Drosophila melanogaster*. *Science*. 2003;302:1727–1736. doi: 10.1126/science.1090289.

Giuliani C, Consonni G, Gavazzi G, Colombo M, Dolfini S. Programmed cell death during embryogenesis in maize. *Ann Bot.* 2002;90:287–292. doi: 10.1093/aob/mcf173.

Gleba Y., Klimyuk V., Marillonnet S. Magniflection—A new platform for expressing recombinant vaccines in plants. *Vaccine*. 2005;23:2042–2048.

Glover, H. Results of Stabilized Rice Bran Evaluation. Turris Phil. Inc. Research & Development-Quality Assurance. December 2007.

Gnanasambandam, R. and N.S. Hetiarachchy, 1995. Protein concentrates from unstabilized and

stabilized rice bran: Preparation and properties. *J. Food Sci.*, 60: 1066-1069 & 74.

Godet, M.C., Tran, V. & Delagw, M.M. (1993). Molecular modelling of the specific interactions in amylose complexation by fatty acids. *International Journal of Biological Macromolecules*, 15, 11–16.

Golde WT. GPRL. A rapid, simple, and humane method for submandibular bleeding of mice using a lancet. *Lab Anim (NY)* 2005;34:39–43.

Goldstein, I.J., R.C. Hughes, M. Monsigny, T. Osawa and N. Sharon, 1980. What should be called a lectin? *Nature*, 285: 66.

Gomi, Y., Fukuoka, M., Mihori, T. & Watanabe, H. (1998). The rate of starch gelatinization as observed byPFG-NMR measurement of water diffusivity in rice starch/water mixtures. *Journal of Food Engineering*, 36, 359–369.

Gomord V., Fitchette A.C., Menu-Bouaouiche L., Saint-Jore-Dupas C., Plasson C., Michaud D., Faye L. Plant-specific glycosylation patterns in the context of therapeutic protein production. *Plant Biotechnol. J.* 2010;8:564–587.

Gonipeta B, Parvataneni S, Paruchuri P, Gangur V. Long-term characteristics of hazelnut allergy in an adjuvant-free mouse model. *Int Arch Allergy Immunol.* 2010;152:219–225. doi: 10.1159/000283028.

Goodard, M.S. Young G: Marcus, R (1984) effect of amylase content on insulin and glucose responses to ingested rice A.M.J. *clin Nutrition* pg, 39, 388-392.

Goodfellow, B.J. & Wilson, R.H. (1990). A Fourier transformIR study of the gelation of amylose and amylopectin. *Biopolymers*, 30, 1183–1189.

Goodman M. Market watch: Sales of biologics to show robust growth through to 2013. *Nat. Rev. Drug Discov.* 2009;8:837.

Gora-Sochacka A., Redkiewicz P., Napiorkowska B., Gaganidze D., Brodzik R., Sirko A. Recombinant mouse granulocyte-macrophage colony-stimulating factor is glycosylated in transgenic tobacco and maintains its biological activity. *J. Interferon Cytokine Res.* 2010;30:135–142.

Grandbastien M.A., Lucas H., Morel J.B., Mhiri C., Vernhettes S., Casacuberta J.M. The expression of the tobacco Tnt1 retrotransposon is linked to plant defense responses. *Genetica.* 1997;100:241–252.

Greenberg JT. Programmed cell death in plant-pathogen interactions. *Ann Rev Plant Physiol Plant Mol Biol.* 1997;48:525–545. doi: 10.1146/annurev.arplant.48.1.525.

Greenberg JT, Yao N. The role and regulation of programmed cell death in plant-pathogen interactions. *Cell Microbiol.* 2004;6:201–211. doi: 10.1111/j.1462-5822.2004.00361.x.

Greenham T., Altosaar I. Molecular strategies to engineer transgenic rice seed compartments for large-scale production of plant-made pharmaceuticals. *Methods Mol. Biol.* 2013;956:311–326.



Grover A, Chandramouli A, Agarwal S, Katiyar-Agarwal S, Agarwal M, Sahi C. In: Rice Improvement in the Genomic Era. Datta SK, editor. Hawarth Press USA; 2009. Transgenic rice for tolerance against abiotic stresses; pp. 237–267.

Gu C.M., Pan H.B., Sun Z.W., Qin G.X. Effect of soybean variety on anti-nutritional factors content, and growth performance and nutrients metabolism in rat. *Int. J. Mol. Sci.* 2010;11:1048–1056.

Gudmundsson, G. (1994). Retrogradation of starch and the role of its components. *Thermochimica Acta*, 246, 329–341.

Gudmundsson, M. & Eliasson, A.-C. (1992). Comparison of thermal and viscoelastic properties of four waxy starches and the effect of added surfactant. *Starch/Stärke*, 44, 379–385.

Guldener U, Munsterkotter M, Oesterheld M, Pagel P, Ruepp A, Mewes HW, Stumpflen V. MPact: the MIPS protein interaction resource on yeast. *Nucleic Acids Res.* 2006;34:D436–441. doi: 10.1093/nar/gkj003.

Gunawardena AH, Greenwood JS, Dengler NG. Programmed cell death remodels lace plant leaf shape during development. *Plant Cell.* 2004;16:60–73. doi: 10.1105/tpc.016188.

Guo C.M., Shi Y.H., Le G.W. Effect of dietary protein level and origin on the redox status in the digestive tract of mice. *Int. J. Mol. Sci.* 2008;9:464–475.

Guraya, H.S., Kadan, R.S. & Champagne, E.T. (1997). Effect of rice starch-lipid complexes on in vitro digestibility, complexing index, and viscosity. *Cereal Chemistry*, 74, 561–565.

Gustafsson C., Govindarajan S., Minshull J. Codon bias and heterologous protein expression. *Trends Biotechnol.* 2004;22:346–353.

H. El-bay A. W 1975 effect of parboiling on the physical and chemical properties of rice geteride mech Brot vol. 29 pg. 230-233.

H. Rivero, J. Mario, H. Raquel, F. Lorena, V. Liliana, and D. Elena, 2007, Concentration of As, Ca, Cd, Cr, Cu, Fe, Hg, K, Mg, Mo, Na, N, Pb and Zn in Uruguayan rice, determined by AAS, Atomic Spectroscopy, 27(2), 48-55.

Hamada A., Yamaguchi K., Ohnishi N., Harada M., Nikumaru S., Honda H. High-level production of yeast (*Schwanniomyces occidentalis*) phytase in transgenic rice plants by a combination of signal sequence and codon modification of the phytase gene. *Plant. Biotechnol. J.* 2005;3:43–55.

Hamada, J.S., 2000. Ultrafiltration of partially hydrolyzed rice bran protein to recover value-added product. *J. Am. Oil Chem. Soc.*, 77: 779-784.

Hamada, J.S., 1997. Characterization of protein fractions of rice bran to devise effective methods of protein solubilization. *Cereal Chem.*, 74: 662-668.

Hamaker, B.R. & Griffin, V.K. (1990). Changing the viscoelastic properties of cooked rice through protein

disruption. *Cereal Chemistry*, 67, 261–264.

Hamaker, B.R. & Griffin, V.K. (1993). Effect of disulfide bound-containing protein on rice starch gelatinization and pasting. *Cereal Chemistry*, 70, 377–380.

Hamaker, B.R., Griffin, V.K. & Moldenhauer, K.A.K. (1991). Potential influence of a starch granule-associated protein on cooked rice stickiness. *Journal of Food Science*, 356, 1327–1330.

Han, B.S., C.B. Park, N. Takasuka, A. Naito, K. Sekine, E. Nomura, H. Taniguchi, T. Tsuno and H. Tsuda, 2001. A ferulic acid derivative, ethyl 3-(4-geranyloxy-3-methoxyphenyl)-2-propenoate, as a new candidate chemopreventive agent for colon carcinogenesis in the rat. *Jpn. J. Cancer Res.*, 92: 404-409.

Han F, Chen H, Li X, Yang M, Liu G, Shen S. A comparative proteomic analysis of rice seedlings under various high-temperature stresses. *Biochim Biophys Acta*. 2009;1794(11):1625–1634.

Han, X.Z. & Hamaker, B.R. (2001). Amylopectin fine structure and rice starch paste breakdown. *Journal of Cereal Science*, 34, 279–284.

Hargrove, K.L.Jr., 1994. Processing and utilization of rice bran in the United States. In: W. E. Marshall and J. I. Wadsworth (Eds.), *Rice science and technology*. New York: Marcel Dekker Inc., pp: 381-404.

Harris MA, Clark J, Ireland A, Lomax J, Ashburner M, Foulger R, Eilbeck K, Lewis S, Marshall B, Mungall C. et al. The Gene Ontology (GO) database and informatics resource. *Nucleic Acids Res*. 2004;32:D258–261. doi: 10.1093/nar/gkh036.

Hartman JW, Tang JE, Wilkinson SB, Tarnopolsky MA, Lawrence RL, Fullerton AV, Phillips SM: Consumption of fat-free fluid milk after resistance exercise promotes greater lean mass accretion than does consumption of soy or carbohydrate in young, novice, male weightlifters. *Am J Clin Nutr* 2007, 86:373-381.

Hatsugai N, Iwasaki S, Tamura K, Kondo M, Fuji K, Ogasawara K, Nishimura M, Hara-Nishimura I. A novel membrane fusion-mediated plant immunity against bacterial pathogens. *Genes Dev*. 2009;23:2496–2506. doi: 10.1101/gad.1825209.

Haug, W. and H.J. Lantzesch, 1983. Sensitive method for the rapid determination of phytate in cereals and cereal products. *J. Sci. Food Agric.*, 34: 1423-1426.

Haumann, B. F. 1989. Rice bran linked to lower cholesterol. *J. Am. Oil Chemists Soc*. 66:615-618.

Hayakawa, T., Seo, S.W. & Igaue, I. (1980). Electron microscopic observation of rice grain. I. Morphology of rice starch. *Journal of the Japanese Society of Starch Science*, 27, 173–179.

He F, Zhang Y, Chen H, Zhang Z, Peng YL. The prediction of protein-protein interaction networks in rice blast fungus. *BMC Genomics*. 2008;9:519. doi: 10.1186/1471-2164-9-519.

- He H, Li J. Proteomic analysis of phosphoproteins regulated by abscisic acid in rice leaves. *Biochem Bioph Res Co.* 2008;371(4):883–888. doi: 10.1016/j.bbrc.2008.05.001.
- He Y., Ning T., Xie T., Qiu Q., Zhang L., Sun Y., Jiang D., Fu K., Yin F., Zhang W., et al. Large-scale production of functional human serum albumin from transgenic rice seeds. *Proc. Natl. Acad. Sci. USA.* 2011;108:19078–19083.
- Health Benefits of Rice Bran: a Review. *J. Nutr.Food Sci.* 2011. 1:3. Nagendra Prasad.
- Heber D: An integrative view of obesity. *Am J Clin Nutr* 2010, 91(suppl):280S-283S.
- Hegsted M., M.M. Windhauser, S. K. Morris, and S. B. Lester. 1993. Stabilized rice bran and oat bran lower cholesterol in humans. *Nutr. Res.* 13:387-398.
- Hellwig S., Drossard J., Twyman R.M., Fischer R. Plant cell cultures for the production of recombinant proteins. *Nat. Biotechnol.* 2004;22:1415–1422.
- Helm, R.M. and A.W. Burks, 1996. Hypoallergenicity of rice protein. *Cereal Foods World*, 41: 839-843.
- Helm RM, Burks AW. Animal models of food allergy. *Curr Opin Allergy Clin Immunol.* 2002;2:541–546. doi: 10.1097/00130832-200212000-00011.
- Hemavathy, J. & Prabhakar, J.V. (1987). Lipid composition of rice (*Oryza sativa* L.) bran. *Journal of the American Oil Chemists' Society*, 64, 1016–1019.
- Hermjakob H, Montecchi-Palazzi L, Lewington C, Mudali S, Kerrien S, Orchard S, Vingron M, Roechert B, Roepstorff P, Valencia A. et al. IntAct: an open source molecular interaction database. *Nucleic Acids Res.* 2004;32:D452–455. doi: 10.1093/nar/gkh052.
- Hernández-Infante, M., V. Sousa, I. Montalvo and E. Tena, 1998. Impact of microwave heating on hemagglutinins, trypsin inhibitors and protein quality of selected legume seeds. *Plant Foods Hum. Nutr.*, 52: 199-208.
- Hiatt A., Cafferkey R., Bowdish K. Production of antibodies in transgenic plants. *Nature.* 1989;342:76–78.
- Hibi, Y. (1998). Roles of water-soluble and water-insoluble carbohydrates in the gelatinization and retrogradation of rice starch. *Starch/Starke*, 50, 474–478.
- Hiei Y, Ohta S, Komari T, Kumashiro T. Efficient transformation of rice (*Oryza sativa* L.) mediated by *Agrobacterium* and sequence analysis of the boundaries of the T-DNA. *Plant J.* 1994;6:271–282. doi: 10.1046/j.1365-313X.1994.6020271.x.
- Hinton, J.J.C. & Shaw, B. (1954). The distribution of nicotinic acid in the rice grain. *British Journal of Nutrition*, 8, 65–71.
- Hiwasa-Tanase K., Nyarubona M., Hirai T., Kato K., Ichikawa T., Ezura H. High-level accumulation of recombinant miraculin protein in transgenic tomatoes expressing a synthetic miraculin gene with

optimized codon usage terminated by the native miraculin terminator. *Plant. Cell Rep.* 2011;30:113–124.

Hizukuri, S. (1985). Relationship between the distribution of the chain length of amylopectin and crystalline structure of starch granule. *Carbohydrate Research*, 141, 295–306.

Hizukuri, S., Kaneko, T. & Takeda, Y. (1983a). Measurement of the chain length of crystalline polymorphism of starch granule. *Biochimica et Biophysica Acta*, 760, 188–191.

Hizukuri, S., Shirasaka, K. & Juliano, B.O. (1983b). Phosphorus and amylose branching in rice starch granules. *Starch/Stärke*, 35, 348–350.

Hizukuri, S., Takeda, Y., Maruta, N. & Juliano, B.O. (1989). Molecular structures of rice starch. *Carbohydrate Research*, 189, 227–235.

Hogan, J.T., Normand, F.L., Deobald, H.J., Mottern, H.H., Lynn, L. & Hunnell, J.W. (1968). Production of high-protein rice flour. *Rice Journal*, 71, 5–6.

Hollak C.E. An evidence-based review of the potential benefits of taliglucerase alfa in the treatment of patients with Gaucher disease. *Core Evid.* 2012;7:15–20.

Hood E.E., Witcher D.R., Maddock S., Meyer T., Baszczynski C., Bailey M., Flynn P., Register J., Marshall L., Bon D., et al. Commercial production of avidin from transgenic maize: Characterization of transformant, production, processing, extraction and purification. *Mol. Breed.* 1997;3:291–306.

Hoogenkamp, H.W.A.P (June 2008). Is Rice Bran Isolate: The New Soy Protein? *Asia Pacific Food Industry*.

Hoogenkamp, Henk. Rice Bran Isolate: The alternative for soy protein? *Poultry Processing*, April 2008, 20-22.

Hoogenkamp, Henk. Building a Better Hybrid Burger. *Fleischwirtschaft International*, Frankfurt Germany. 1/2013, 2/2013. Hoogenkamp, Henk. GMO: For Profit and Humanity, *World of Ingredients*. Feb. 2013.

Hoogenkamp, Henk. Building a Hybrid Burger, *Fleischwirtschaft International* 1/2013, 2/2013.

Hoogenkamp, Henk W. *Milk Protein: The complete guide to meat, poultry and seafood*. 1989. DMV Campina bv. Veghel, The Netherlands.

Hoogenkamp, Henk W. *Vegetable Protein: Interactive Technology and Marketing for Meat, Poultry & Lifestyle Foods*. DuPont Protein Technologies Inc. (1998).

Hoogenkamp, Henk W. *Soy protein and formulated meat products*. ISBN: 085199 864. [www.cabi-publishing.org](http://www.cabi-publishing.org).

- Hoogenkamp, Henk W. Proteina de soja y formulas para productos carnicos. ISBN: 978 84200 1101 1. [www.editorialacribia.com](http://www.editorialacribia.com).
- Hoogenkamp, Henk. Vegetable Protein Marketing. ISBN-13: 978-1466258631.
- Hoogenkamp, Henk. Rice Protein. ISBN 978-1477468142. August 2012.
- Horigane, A.K., Toyoshima, H., Hemmi, H., Engelaar, W.M.H.G., Okubo, A. & Nagata, T. (1999). Internal hollows in cooked rice grains (*Oryza sativa* cv. Koshihikari) observed by NMR micro imaging. *Journal of Food Science*, 64, 1–5.
- Horn M.E., Woodard S.L., Howard J.A. Plant molecular farming: Systems and products. *Plant Cell Rep.* 2004;22:711–720.
- Houghton JT, Ding Y, Griggs DJ, Nogue M, van der Linden PJ, Dai X, Maskell K, Johnson CA. Climate change 2001: The scientific basis. Cambridge UK: Cambridge University Press; 2001. pp. 1–20.
- Houston, D. and A. Mohammad. 1970. Purification and partial characterization of a major globulin from rice endosperm. *Cereal Chem.*, 47: 5-11.
- Houston, D.F. (1967). High protein flour can be made from all types of milled rice. *Rice Journal*, 70, 12–15.
- Houston, D., T. Iwasaki, A. Mohammad and L. Chen. 1968. Radial distribution of protein by solubility classes in the milled rice kernel. *J. Agri. Food Chem.*, 16: 720-724.
- Hsieh K.Y., Hsu C.I., Lin J.Y., Tsai C.C., Lin R.H. Oral administration of an edible-mushroom-derived protein inhibits the development of food-allergic reactions in mice. *Clin. Exp. Allergy*. 2003;33:1595–1602.
- Hsu, S., Lu, S. & Huang, C. (2000). Viscoelastic changes of rice starch suspensions during gelatinization. *Journal of Food Science*, 65, 215–220.
- Hu E.A., Pan A., Malik V., Sun Q. White rice consumption and risk of type 2 diabetes: Meta-analysis and systematic review. *Br. Med. J.* 2012;344 doi: 10.1136/bmj.e1454.
- Hu F.B., van Dam R.M., Liu S. Diet and risk of Type II diabetes: The role of types of fat and carbohydrate. *Diabetologia*. 2001;44:805–817.
- Hu, W., J.H. Wells, T.S. Shin and J.S. Godber, 1996. Comparison of isopropanol and hexane for extraction of vitamin E and oryzanols from stabilized rice bran. *J. Am. Oil Chem. Soc.*, 73: 1653-1656.
- Huang J., Sutliff T.D., Wu L., Nandi S., Bengte K., Terashima M., Ralston A.H., Drohan W., Huang N., Rodriguez R.L. Expression and purification of functional human alpha-1-antitrypsin from cultured plant cells. *Biotechnol. Prog.* 2001;17:126–133.

Huang L.F., Liu Y.K., Lu C.A., Hsieh S.L., Yu S.M. Production of human serum albumin by sugar starvation induced promoter and rice cell culture. *Transgenic Res.* 2005;14:569–581.

Huang T.K., McDonald K.A. Bioreactor systems for in vitro production of foreign proteins using plant cell cultures. *Biotechnol. Adv.* 2012;30:398–409.

Huang TW, Lin CY, Kao CY. Reconstruction of human protein interolog network using evolutionary conserved network. *BMC Bioinformatics.* 2007;8:152. doi: 10.1186/1471-2105-8-152.

Huang HD, Lee TY, Tzeng SW, Horng JT. KinasePhos: a web tool for identifying protein kinase-specific phosphorylation sites. *Nucleic Acids Res.* 2005;33:W226–W229. doi: 10.1093/nar/gki471.

Huang T.-K., McDonald K.A. Bioreactor engineering for recombinant protein production in plant cell suspension cultures. *Biochem. Bioeng. J.* 2009;45:168–184.

Huebner, F., J. Bietz, B. Webb and B. Juliano. 1990. Rice cultivar identification by high-performance liquid chromatography of endosperm proteins. *Cereal Chem.*, 67: 129-135.

Human Colon and Liver Cancer Cell Proliferation Inhibition by Peptide Hydrolysates Derived from Heat-Stabilized Defatted Rice Bran. *Journal of Agricultural and Food Chemistry*, 2008, 56, 11643-11647.

Hundemer, J. K., S. P. Nabar, B. J. Shriver, and L. P. Forman. 1991. Dietary fiber Sources lower blood cholesterol in C57BL/6 mice. *J. Nutr.* 121:1360-1365.

Iai V.M.F. Lu, S and Lii, (2000) molecular characteristics influencing retrogradation kinetic of rice amylopectins *cereal chemistry* vol. 77 pg. 315-319.

Induhara Swamy, Y.M Sowbhaya. C.M. and Bhattacharya, K.R. (1978) changes in the physicochemical properties of rice with ageing. *Journal science food Agric* vol. 29 pg. 627-639.

Ihekoronye A.I. and Ngoddy (1985). *Integrated food science technology for the tropics* mamillian publisher. Ltd., London, pg. 255.

Ihekoronye A.I. and Ngoddy (1985). *Integrated food science technology for the tropics* mamillian publisher. Ltd., London, pg. 255.

Ihekoronye A.I. and Ngoddy (1985). *Integrated food science technology for the tropics* mamillian publisher. Ltd., London, pg. 255.

Ikezawa Z, Tsubaki K, Osuna H, Shimada T, Moteki K, Sugiyama H, Katumata K, Anzai H, Amano S. Usefulness of hypoallergenic rice (AFT-R 1) and analysis of the salt insoluble rice allergen molecule. *Arerugi.* 1999;48:40–49.

Ikezawa Z, Miyakawa K, Komatsu H, Suga C, Miyakawa J, Sugiyama A, Sasaki T, Nakajima H, Hirai Y, Suzuki Y. A probable involvement of rice allergy in severe type of atopic dermatitis in Japan. *Acta Derm Venereol Suppl (Stockh)* 1992;176:103–107.

Ikezawa Z, Ikebe T, Ogura H, Odajima H, Kurosaka F, Komatu H, et al. Mass trial of hypoallergenic rice (HRS-1) produced by enzymatic digestion in atopic dermatitis with suspected rice allergy. *Acta Derm Venereol Suppl (Stockh)* 1992;176:108–112.

Induhara Swamy, Y.M Sowbhaya. C.M. and Bhattacharya, K.R. (1978) changes in the physiochemical prosperities of rice with ageing. *Journal science food Agric* vol. 29 pg. 627-639.

Ingelman-Sundberg M. Genetic polymorphisms of cytochrome P450 2D6 (CYP2D6): Clinical consequences, evolutionary aspects and functional diversity. *Pharmacogenomics J.* 2005;5:6–13.

International Rice Commission. (1998). 19th Session September (1998). Cairo: FAO.

IRGSP. The map-based sequence of the rice genome. *Nature.* 2005;436:793–800.

Iritani N, Hosomi H, Fukuda H, Tada K, Ikeda H: Soybean protein suppresses hepatic lipogenic enzyme gene expression in Wistar fatty rats. *J Nutr* 1996, 126:380-388. Induhara Swamy, Y.M Sowbhaya. C.M. and Bhattacharya, K.R. (1978) changes in the physiochemical prosperities of rice with ageing.

Ishibashi, G., and M. Yamamoto. 1980. Effect of rice wax, garlic, and ginkgo seed component on plasma and liver cholesterol levels in rats. *Kynshi Women's Univ.* 16:115-127.

Isono, H., Ohtsubo, K., Iwasaki, T. & Yamazaki, A. (1994). Eating quality of domestic and foreign rices of various varieties and characteristics. *Journal of the Japanese Society for Food Science and Technology-Nippon Shokuhin Kogyo Gakkaishi*, 41, 485–492.

Ito M, Kato T, Matsuda T. Rice allergenic proteins, 14-16 kDa albumin and alpha-globulin, remain insoluble in rice grains recovered from rice miso (rice-containing fermented soybean paste) *Biosci Biotechnol Biochem.* 2005;69:1137–1144.

Ito, S., Sato, S. & Fujino, Y. (1979). Internal lipid in rice starch. *Starch/Stärke*, 31, 217–221.

Itoh, T., T. Tamura, and T. Matsumoto. 1973. Sterol composition of 19 vegetable oils. *J. Am. Oil Chemists Soc.* 50:122-125.

Izumi H, Sugiyama M, Matsuda T, Nakamura R. Structural characterization of the 16-kDa allergen, RA17, in rice seeds. Prediction of the secondary structure and identification of intramolecular disulfide bridges. *Biosci Biotechnol Biochem.* 1999;63:2059–2063.

Jacobson, M.R., Obanni, M. & BeMiller, J.M. (1997). Retrogradation of starches from different botanical sources. *Cereal Chemistry*, 74, 511–518.

Jain M, Nijhawan A, Tyagi AK, Khurana JP. Validation of housekeeping genes as internal control for studying gene expression in rice by quantitative real-time PCR. *Biochem Biophys Res Commun.* 2006;345:646–651. doi: 10.1016/j.bbrc.2006.04.140.

- Jang H, Chi Y, Park S, Lee S, Lee J, Park J, Moon J, Lee Y, Kim S, Lee K, Lee S. Structural and functional regulation of eukaryotic 2-Cys peroxiredoxins including the plant ones in cellular defense signaling mechanisms against oxidative stress. *Physiol Plantarum*. 2006;126:549–559. doi: 10.1111/j.1399-3054.2006.00685.x.
- Jaquinod M, Villiers F, Kieffer-Jaquinod S, Hugouvieux V, Bruley C, Garin J, Bourguignon J. A proteomics dissection of *Arabidopsis thaliana* vacuoles isolated from cell culture. *Mol Cell Proteomics*. 2007;6:394–412.
- Jane, J., Chen, Y.Y., Lee, L.F., et al. (1999). Effects of amylopectin branch chain length and amylose content on the gelatinization and pasting properties of starch. *Cereal Chemistry*, 76, 629–637.
- Jansen R, Yu H, Greenbaum D, Kluger Y, Krogan NJ, Chung S, Emili A, Snyder M, Greenblatt JF, Gerstein M. A Bayesian networks approach for predicting protein-protein interactions from genomic data. *Science*. 2003;302:449–453. doi: 10.1126/science.1087361.
- Jayadeep A., Malleshi N.G. Nutrients, composition of tocotrienols, tocopherols, and  $\gamma$ -oryzanol, and antioxidant activity in brown rice before and after biotransformation. *CyTA J. Food*. 2011;9:82–87.
- Jenkins D.J.A., Kendall C.W.C., Augustin L.S.A., Franceschi S., Hamidi M., Marchie A., Jenkins A.L., Axelsen M. Glycemic index: Overview of implications in health and disease. *Am. J. Clin. Nutr.* 2002;76:266S–273S.
- Jin C., Altmann F., Strasser R., Mach L., Schahs M., Kunert R., Rademacher T., Glossl J., Steinkellner H. A plant-derived human monoclonal antibody induces an anti-carbohydrate immune response in rabbits. *Glycobiology*. 2008;18:235–241.
- Jin M, Diaz PT, Bourgeois T, Eng C, Marsh CB, Wu HM. Two-dimensional gel proteome reference map of blood monocytes. *Proteome sci*. 2006;4:16. doi: 10.1186/1477-5956-4-16. Johnson I. T., J. M. Gee, and J. C. Brown. 1989. A comparison of rice bran, wheat bran and cellulose as sources of dietary fibre in the rat. *Food Sci. Nutr.* 42:153-163.
- Jones JDG, Dangl JL. The plant immune system. *Nature*. 2006;444:323–329. doi: 10.1038/nature05286.
- Jones SM, Magnolfi CF, Cooke SK, Sampson HA. Immunologic cross-reactivity among cereal grains and grasses in children with food hypersensitivity. *J Allergy Clin Immunol*. 1995;96:341–351.
- Journal science food Agric vol. 29 pg. 627-639.
- Ju, Z., N. Hettiarachchy and N. Rath. 2001. Extraction, denaturation and hydrophobic properties of rice flour proteins. *J. Food Sci.*, 66: 229-232.
- Juliano, B.O. (1979). The chemical basis of rice grain quality. *Proceedings of a Workshop on Chemical Aspects of Rice Grain Quality*. Pp. 69–90. Los Ban~ os, Laguna, Philippines: International Rice Research Institute.



- Juliano, B.O. 1972. Physico-chemical properties of starch and protein and their relation to grain quality and nutritional value of rice. Rice Breeding. International Rice Research Institute, Los Banos, Philippines.
- Juliano, B.O. (1983). Lipids in rice and rice processing. In: Lipids in Cereal Technology (edited by P.J. Barnes). Pp. 305–330. London: Academic Press.
- Juliano, B.O. (1985a). Production and utilization of rice. In: Rice Chemistry and Technology (edited by B.O. Juliano). Pp. 1–16. St Paul, Minnesota: American Association of Cereal Chemists.
- Juliano, B.O. (1985b). Criteria and tests for rice grain qualities. In: Rice Chemistry and Technology (edited by B.O. Juliano). Pp. 443–524. St Paul, Minnesota: American Association of Cereal Chemists.
- Juliano, B.O. (1985c). The rice grain and its gross composition. In: Rice Chemistry and Technology (edited by B.O. Juliano). Pp. 17–57. St Paul, Minnesota: American Association of Cereal Chemists.
- Juliano, B.O. (1985d). Polysaccharides, proteins, and lipids of rice. In: Rice Chemistry and Technology (edited by B.O. Juliano). Pp. 59–174. St Paul, Minnesota: American Association of Cereal Chemists.
- Juliano, B.O., Nazareno, M.B. & Ramos, N.B. (1969). Properties of waxy and isogenic non-waxy rices differing in starch gelatinization temperature. *Journal of Agricultural and Food Chemistry*, 17, 1364–1369.
- Juliano, B.O. 1980. Rice: Recent progress in chemistry and nutrition. In: *Cereals for Food and Beverages, Recent Progress in Cereal Chemistry*. (Eds.): G.E. Inglett and L. Munck. Academic Press, New York.
- Juliano, B.O. 1985. Polysaccharides, proteins, and lipids. In: *Rice: chemistry and technology*. (Ed.): B.O. Juliano. AACC, St. Paul, pp. 59-174.
- Jung HW, Hwang BK. The leucine-rich repeat (LRR) protein, CaLRR1, interacts with the hypersensitive induced reaction (HIR) protein, CaHIR1, and suppresses cell death induced by the CaHIR1 protein. *Mol Plant Pathol*. 2007;8:503–514. doi: 10.1111/j.1364-3703.2007.00410.x.
- Jung KH, Dardick C, Bartley LE, Cao P, Phetsom J, Canlas P, Seo YS, Shultz M, Ouyang S, Yuan Q. et al. Refinement of light-responsive transcript lists using rice oligonucleotide arrays: evaluation of gene-redundancy. *PLoS One*. 2008;3:e3337. doi: 10.1371/journal.pone.0003337.
- Jwa NS, Park SG, Park CH, Kim SO, Ahn IP, Park SY, Yoon CH, Lee YH. Cloning and expression of a rice cDNA encoding a Lls1 homologue of maize. *Plant Pathol J*. 2000;16:151–155.
- K. Tanaka, M. Ogawa and Z. Kasai, 1977, The rice scutellumII, *Journal of Cereal Chemistry*, 54, 684-689.
- Kagawa, H., H. Hirano and F. Kikuchi. 1988. Variation of glutelin seed storage protein in rice (*Oryza sativa* L.). *Ikushugaku Zasshi*, 38: 327-332.

- Kahlon, T. S., F. I. Chow, and D. F. Wood. 1999. Cholesterol response and foam cell formation in hamsters fed rice bran, oat bran, and cellulose + soy protein diets with or without added vitamin E. *Cereal Chem.* 76:772-776.
- Kahlon, T.S. and G.E. Smith, 2004. Rice bran: A health promoting ingredient. *Cereal Foods World*, 49: 188-194. Kato N, Iwami K: Resistant protein: its existence and function beneficial to health. *J Nutr Sci Vitaminol* 2002, 48:1-5.
- Kahlon T. S., R. M. Saunders, R. N. Sayre, F. I. Chow, M. M. Chiu, and A. A. Betschart 1990. Influence of rice bran, oat bran, and wheat bran on cholesterol and triglycerides in hamsters. *Cereal Chem.* 67:439-443.
- Kahlon T. S., F. I. Chow, R. N. Sayre, and A. A. Betschart. 1992. Cholesterol lowering in hamsters fed rice bran at various levels, defatted rice bran and rice bran oil. *J. Nutr.* 122:513-519.
- Kahlon T. S., F. I. Chow, B. E. Knuckles, and M. M. Chiu. 1993. Cholesterol lowering effects in hamsters of 3g1ucanenriched barley fraction, dehulled whole barley, rice bran, and oat bran and their combinations. *Cereal Chem.* 70:435-440.
- Kahlon, T. S., R. M. Saunders, R. N. Sayre, F. I. Chow, M. M. Chiu, and A. A. Betschart. 1992. Cholesterol-lowering effects of rice bran and rice bran oil fractions in hypercholesterolemic hamsters. *Cereal Chem.* 69:485-489.
- Kahlon T. S., R. M. Saunders, R. N. Sayre, F. I. Chow, M. M. Chiu, and A. A. Betschart. 1989. Effect of rice bran and oat bran on plasma cholesterol in hamsters. *Cereal Foods World* 34:768 .
- Kahlon T. S., F. I. Chow, M. M. Chiu, C. A. Hudson, and R. N. Sayre. 1996. Cholesterol-lowering by rice bran and rice bran oil unsaponifiable matter in hamsters. *Cereal Chem.* 73:69-74.
- Kahlon, T. S., R.H. Edwards, and F. I. Chow. 1998. Effect of extrusion on hypocholesterolemic properties of rice, oat, corn, and wheat bran diets in hamsters. *Cereal Chem.* 75:897-903. Aoe S., F. Ohta, and Y.
- Kahlon, T. S., and F. I. Chow. 2000. In vitro binding of bile acids by rice bran, oat bran, wheat bran and corn bran. *Cereal Chem.* 77: 518-521.
- Kahlon, T. S., and Woodruff, C. L. 2003. In vitro binding of bile acids by rice bran, oat bran, barley and 3-glucan enriched barley. *Cereal Chem.* 80:260-263.
- Kalichevsky, M.T., Oxford, P.D. & Ring, S.G. (1990). The retrogradation and gelation of amylopectins from various botanical sources. *Carbohydrate Research*, 198, 49–55.
- Kang SW, Changa TS, Lee TH, Kima ES, Yue DY, Rhee SG. Cytosolic peroxiredoxin attenuates the activation of Jnk and p38 but potentiates that of Erk in Hela cells stimulated with tumor necrosis factor- $\alpha$ . *J Biol Chem.* 2004;279(4):2535–2543.
- Kanekatsu M, Saito H, Motohashi K, Hisabori T. The beta subunit of chloroplast ATP synthase (CF<sub>0</sub>CF<sub>1</sub>-ATPase) is phosphorylated by casein kinase II. *Biochem Mol Biol Int.* 1998;46(1):99–105.

- Kanekatsu M, Ezumi A, Nakamura T, Ohtsuki K. Chloroplast Ribonucleoproteins (RNPs) as Phosphate Acceptors for Casein Kinase II: Purification by ssDNA-Cellulose Column Chromatography. *Plant Cell Physiol.* 1995;36(8):1649–1656.
- Kanno T., Naito S., Shimamoto K. Post-transcriptional gene silencing in cultured rice cells. *Plant. Cell. Physiol.* 2000;41:321–326.
- Kapila J., de Rycke R., van Montagu M., Angenon G. Agrobacterium-mediated transient gene expression system for intact leaves. *Plant Sci.* 1997;122:101–108.
- Karkalas, J., Ma, S. & Morrison, W.R. (1995). Some factors determining the thermal properties of amylose inclusion complexes with fatty acids. *Carbohydrate Research*, 268, 233–247.
- Katagiri F, Thilmony R, He SY. In: *The Arabidopsis Book*. Somerville CR, Meyerowitz EM, editor. Vol. 39. Rockville, MD: American Society of Plant Biologists; 2002. The Arabidopsis thaliana-Pseudomonas syringae interaction.
- Katial RK, Grier TJ, Hazelhurst DM, Hershey J, Engler RJ. Deleterious effects of electron beam radiation on allergen extracts. *J Allergy Clin Immunol.* 2002;110:215–219.
- Kato N, Iwami K: Resistant protein: its existence and function beneficial to health. *J Nutr Sci Vitaminol* 2002, 48:1-5.
- Kaufmann H, Bailey JE, Fussenegger M. Use of antibodies for detection of phosphorylated proteins separated by two-dimensional gel electrophoresis. *Proteomics.* 2001;1(2):194–199. doi: 10.1002/1615-9861(200102)1:2<194::AID-PROT194>3.0.CO;2-K.
- Kaul R, Saluja D, Sachar RC. Phosphorylation of small subunit plays a crucial role in the regulation of RuBPCase in moss and spinach. *FEBS.* 1986;209:63–70. doi: 10.1016/0014-5793(86)81085-7.
- Kaundal R, Raghava GP. RSLpred: an integrative system for predicting subcellular localization of rice proteins combining compositional and evolutionary information. *Proteomics.* 2009;9:2324–2342. doi: 10.1002/pmic.200700597.
- Kaur, K. & Singh, N. (2000). Amylose-lipid complex formation during cooking of rice flour. *Food Chemistry*, 71, 511–517.
- Kawamura, Y., M. Muramoto, K.W. Waldron, I.T. Johnson and L.R. Fenwick, 1993. Anti-tumorigenic and immunoactive protein and peptide factors in foodstuffs. 2. Anti-tumorigenic factors in rice bran. *Food Cancer Prev.: Chem. Biol. Aspects.*, pp: 331-401.
- Kawashima, K. & Kiribuchi, T. (1980). Studies on lipid components and heat dependent pasting behaviour of non-waxy and waxy rice starches. *Kaseigaku Zasshi*, 31, 625–628.

Kayashita J., Shimoka I., Nakajoh M., Yamazaki M., Kato N. Consumption of buckwheat protein lowers plasma cholesterol and raises fecal neutral sterols in cholesterol-fed rats because of its low digestibility. *J. Nutr.* 1997;127:1395–1400.

Ke Y, Han G, He H, Li J. Differential regulation of proteins and phosphoproteins in rice under drought stress. *Biochem Bioph Res Co.* 2009;379(1):133–138. doi: 10.1016/j.bbrc.2008.12.067.

Keshava Prasad TS, Goel R, Kandasamy K, Keerthikumar S, Kumar S, Mathivanan S, Telikicherla D, Raju R, Shafreen B, Venugopal A. et al. Human Protein Reference Database--2009 update. *Nucleic Acids Res.* 2009;37:D767–772. doi: 10.1093/nar/gkn892.

Kestin M., R. Moss, C. M. Clifton, and P. J. Nestle]. 1990. Comparative effects of three cereal brans on plasma lipids, blood pressure, and glucose metabolism in mildly hypercholesterolemic men. *Am. J. Clin. Nutr.* 52:661-666.

Khan M, Takasaki H, Komatsu S. Comprehensive phosphoproteome analysis in Rice and identification of phosphoproteins responsive to different hormones/stresses. *J Proteome Res.* 2005;4(5):1592–1599. doi: 10.1021/pr0501160.

Khush G. Productivity improvements in rice. *Nutr Rev.* 2003;61:S114–116. doi: 01301/nr.2003.jun.S114-S116.

Kieran P.M., Malone D.M., MacLoughlin P.F. Effects of hydrodynamic and interfacial forces on plant cell suspension systems. *Adv. Biochem. Eng. Biotechnol.* 2000;67:139–177.

Kilshaw PJ, Heppell LM, Ford JE. Effects of heat treatment of cow's milk and whey on the nutritional quality and antigenic properties. *Arch Dis Child.* 1982;57:842–847.

Kim D.I., Cho G.H., Pedersen H., Chin C.K. A hybrid bioreactor for high density cultivation of plant cell suspensions. *Appl. Microbiol. Biotechnol.* 1991;34:726–729.

Kim H.J., Lee D.H., Kim D.K., Han G.B. The glycosylation and in vivo stability of human granulocyte-macrophage colony-stimulating factor produced in rice cells. *Biol. Pharm. Bull.* 2008;31:290–294.

Kim N.S., Kim T.G., Jang Y.S., Shin Y.J., Kwon T.H., Yang M.S. Amylase gene silencing by RNA interference improves recombinant hGM-CSF production in rice suspension culture. *Plant. Mol. Biol.* 2008;68:369–377.

Kim N.S., Kim T.G., Kim O.H., Ko E.M., Jang Y.S., Jung E.S., Kwon T.H., Yang M.S. Improvement of recombinant hgm-csf production by suppression of cysteine proteinase gene expression using RNA interference in a transgenic rice culture. *Plant. Mol. Biol.* 2008;68:263–275.

Kim, S.K., Hahn, T.R., Lee, Y.H. & D'Appolonia, B.L. (1978). Physicochemical properties of tongil (Indica type) and paldal (Japonica type) rice starch. *Korean Journal of Food Science and Technology*, 10, 157–165.

- Kim T.G., Lee H.J., Jang Y.S., Shin Y.J., Kwon T.H., Yang M.S. Co-expression of proteinase inhibitor enhances recombinant human granulocyte-macrophage colony stimulating factor production in transgenic rice cell suspension culture. *Protein Expr. Purif.* 2008;61:117–121.
- Kim TH, Jung HH, Kim ES, Park JY, Kim KW, Sohn MH, et al. A case of rice allergy caused by thin rice porridge during the weaning period. *Pediatr Allergy Respir Dis.* 2007;17:149–154.
- Kim T.G., Baek M.Y., Lee E.K., Kwon T.H., Yang M.S. Expression of human growth hormone in transgenic rice cell suspension culture. *Plant Cell. Rep.* 2008;27:885–891.
- Kim WY, Kang S, Kim BC, Oh J, Cho S, Bhak J, Choi JS. SynechoNET: integrated protein-protein interaction database of a model cyanobacterium *Synechocystis* sp. PCC 6803. *BMC Bioinformatics.* 2008;9(Suppl 1):S20. doi: 10.1186/1471-2105-9-S1-S20.
- Kimber I, Dearman RJ, Penninks AH, Knippels LM, Buchanan RB, Hammerberg B, Jackson HA, Helm RM. Assessment of protein allergenicity on the basis of immune reactivity: animal models. *Environ Health Perspect.* 2003;111:1125–1130.
- Kishimoto, T., M. Watanabe, T. Mitsui and H. Hori. 1999. Glutelin basic subunits have a mammalian mucin-type Olinked disaccharide side chain. *Arch. Biochem. Biophysics*, 370: 271-277.
- Kishine M., Suzuki K., Nakamura S., Ohtsubo K. Grain qualities and their genetic derivation of 7 new rice for Africa (NERICA) varieties. *J. Agric. Food Chem.* 2008;56:4605–4610.
- Kitahara, K., Suganuma, T. & Nagahama, T. (1996). Susceptibility of amylose-lipid complexes to hydrolysis by glucoamylase from *Rhizopus niveus*. *Cereal Chemistry*, 73, 428–432.
- Kitahara, K., Tanaka, T., Suganuma, T. & Nagahama, T. (1997). Release of bound lipids in cereal starches upon hydrolysis by glucoamylase. *Cereal Chemistry*, 74, 1–6.
- Koch E, Slusarenko A. Arabidopsis is susceptible to infection by a downy mildew fungus. *Plant Cell.* 1990;2:437–445. doi: 10.1105/tpc.2.5.437.
- Kraemer WJ, Ratamess NA, Volek JS, Hakkinen K, Rubin MR, French DN, Gomez AL, McGuigan MR, Scheett TP, Newton RU, et al.: The effects of amino acid supplementation on hormonal responses to resistance training overreaching. *Metabolism* 2006, 55:282-291.
- Kraemer WJ, Hatfield DL, Volek JS, Fragala MS, Vingren JL, Anderson JM, Spiering BA, Thomas GA, Ho JY, Quann EE, et al.: Effects of amino acids supplement on physiological adaptations to resistance training. *Medicine and science in sports and exercise* 2009, 41:1111-1121.
- Krishnan, H. and J. White. 1995. Morphometric analysis of rice seed protein bodies. Implication for a significant contribution of prolamine to the total protein content of rice endosperm. *Plant Physiol.*, 109: 1491-1495.

Krogan NJ, Cagney G, Yu H, Zhong G, Guo X, Ignatchenko A, Li J, Pu S, Datta N, Tikuisis AP. et al. Global landscape of protein complexes in the yeast *Saccharomyces cerevisiae*. *Nature*. 2006;440:637–643. doi: 10.1038/nature04670.

Kubota M., Sato Y., Masumura T., Kumagai T., Watanabe R., Fujimura S., Kadowali M. Improvement in the in vitro digestibility of rice protein by alkali extraction is due to structural changes in prolamin/protein body-I particle. *Biosci. Biotechnol. Biochem.* 2010;74:614–619.

Kugimiya, M., Donovan, J.W. & Wong, R.Y. (1980). Phase transitions of amylose-lipid complexes in starches: a calorimetric study. *Starch/Stärke*, 32, 265–270.

Kuiper PJC. In: Ann NY Acad Sci. Csermely P, editor. Vol. 851. 1998. Adaptation mechanism of green plants to environmental stress. The role of plant sterols and the phosphatidyl linolenoyl cascade in the functioning of plants and the response of plants to global climate change; pp. 209–215. *Stress of Life: From Molecules to Man*.

Kum, J.S., H.W. Yoo and J.D. Park. 2010. Strategies for the development of rice processing technologies and industries. In: 2010 KREI Agricultural Outlooks. Korea Rural Economic Institute (KREI), Korea. pp. 603-633.

Kumagai T., Kawamura H., Fuse T., Watanabe T., Sato Y., Masumura T., Watanabe R., Kadowaki M. Production of rice protein by alkaline extraction improves its digestibility. *J. Nutr. Sci. Vitaminol.* 2006;52:467–472.

Kumar R, Srivastava P, Kumari D, Fakhr H, Sridhara S, Arora N, Gaur SN, Singh BP. Rice (*Oryza sativa*) allergy in rhinitis and asthma patients: a clinico-immunological study. *Immunobiology.* 2007;212:141–147. doi: 10.1016/j.imbio.2006.11.006.

Kuo-Chen Chou, Hong-Bin Shen. Large-scale plant protein subcellular location prediction. *Journal of Cellular Biochemistry.* 2007;100:665–678. doi: 10.1002/jcb.21096.

Kusnadi A.R., Nikolov Z.L., Howard J.A. Production of recombinant proteins in transgenic plants: Practical considerations. *Biotechnol. Bioeng.* 1997;56:473–484.

Kwon, D.H., J.H. Seung, S.H. Han and D.G. Park. 2012. Korea Rice Supply-Demand Outlook. In: 2012 KREI Agricultural Outlooks. Korea Rural Economic Institute (KERI), Korea. pp. 484-508.

Kwon J.Y., Lee K.H., Cheon S.H., Ryu H.N., Kim S.J., Kim D.I. Adsorptive loss of secreted recombinant proteins in transgenic rice cell suspension cultures. *Plant. Cell Rep.* 2012;31:551–560.

Kumar R, Srivastava P, Kumari D, Fakhr H, Sridhara S, Arora N, et al. Rice (*Oryza sativa*) allergy in rhinitis and asthma patients: a clinico-immunological study. *Immunobiology.* 2007;212:141–147.

Laemmli, U. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, 227: 680-685.

- Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*. 1970;227:680–685.
- Lai, H.M. (2001). Effects of hydrothermal treatment on the physicochemical properties of pregelatinized rice flour. *Food Chemistry*, 72, 455–463.
- Lai, V.M.F. & Lii, C.Y. (1999). Effects of modulated differential scanning calorimetry (MDSC) variables on thermodynamic and kinetic characteristics during gelatinization of waxy rice starch. *Cereal Chemistry*, 76, 519–525.
- Lai, V.M.F., Lu, S. & Lii, C. (2000). Molecular characteristics influencing retrogradation kinetics of rice amylopectins. *Cereal Chemistry*, 77, 272–278.
- Lai, V.M.F., Shen, M.C., Yeh, A.I., Juliano, B.O. & Lii, C.Y. (2001). Molecular and gelatinization properties of rice starches from IR24 and Sinandomeng cultivars. *Cereal Chemistry*, 78, 596–602.
- Lakoucheva LM, Radivojac P, Brown CJ, O'Connor TR, Sikes JG, Obradovic Z, Dunker AK. The importance of intrinsic disorder for protein phosphorylation. *Nucleic Acids Res*. 2004;32(3):1037–1049. doi: 10.1093/nar/gkh253.
- Lam SK, Siu CL, Hillmer S, Jang S, An G, Robinson DG, Jiang L. Rice SCAMP1 defines clathrin-coated, trans-Golgi-located tubular-vesicular structures as an early endosome in tobacco BY-2 cells. *Plant Cell*. 2007;19:296–319. doi: 10.1105/tpc.106.045708.
- Langdon T., Seago C., Mende M., Leggett M., Thomas H., Forster J.W., Jones R.N., Jenkins G. Retrotransposon evolution in diverse plant genomes. *Genetics*. 2000;156:313–325.
- Larson, B.L., Gilles, K.A. & Jenness, R. (1953). Amperometric method for determining the sorption of iodine by starch. *Analytical Chemistry*, 25, 802–804.
- La'sztity, R. (1999). The chemistry of rice. In: *Cereal Chemistry* (edited by R. La'sztity). Pp. 267–290. Budapest: Akadé'miai Kiado'.
- Lawn R.M., Adelman J., Bock S.C., Franke A.E., Houck C.M., Najarian R.C., Seeburg P.H., Wion K.L. The sequence of human serum albumin cDNA and its expression in *E. coli*. *Nucleic Acids Res*. 1981;9:6103–6114.
- Layman DK: Role of leucine in protein metabolism during exercise and recovery. *Can J Appl Physiol* 2002, 27:646-663.
- L'hocine, L., J.I. Boye and Y. Arcand, 2006. Composition and functional properties of soy protein isolates prepared using alternative defatting and extraction procedures. *J. Food Sci.*, 71: C137-C145.
- Lee C.R., Pieper J.A., Frye R.F., Hinderliter A.L., Blaisdell J.A., Goldstein J.A. Tolbutamide, flurbiprofen, and losartan as probes of CYP2C9 activity in humans. *J. Clin. Pharmacol*. 2003;43:84–91.

Lee D, Ahsan NB, Lee S, Kang K, Bahk J, Lee I, Lee B. A proteomic approach in analyzing heat-responsive proteins in rice leaves. *Proteomics*. 2007;7(18):3369–3383. doi: 10.1002/pmic.200700266.

Lee K.T., Chen S.C., Chiang B.L., Yamakawa T. Heat-inducible production of beta-glucuronidase in tobacco hairy root cultures. *Appl. Microbiol. Biotechnol.* 2007;73:1047–1053.

Lee S.J., Park C.I., Park M.Y., Jung H.S., Ryu W.S., Lim S.M., Tan H.K., Kwon T.H., Yang M.S., Kim D.I. Production and characterization of human CTLA4Ig expressed in transgenic rice cell suspension cultures. *Protein Expr. Purif.* 2007;51:293–302.

Lee G.-H., Sohn S.-H., Park E.-Y., Park Y.-D. Aberrant promoter methylation occurred from multicopy transgene and SU(VAR)3–9 homolog 9 (SUVH9) gene in transgenic *Nicotiana benthamiana*. *Funct. Plant Biol.* 2012;39:764–773.

Lee KY, Lee SY, Jeong BJ, Kim KE, Kim DS. Cross-reactivity among four cereals: rice, barley, wheat flour and buckwheat flour. *Korean J Allergy*. 1993;13:65–74.

Lee SY, Jeoung BJ, Kim DS, Kim KE, Lee KY. Detection of specific IgE and IgG4 to rice and its significance in atopic children. *Pediatr Allergy Respir Dis.* 1993;3:74–82.

Lee Y.T., Lee S.S., Sun H.L., Lu K.H., Ku M.S., Sheu J.N., Ko J.L., Lue K.H. Effect of the fungal immunomodulatory protein FIP-fve on airway inflammation and cytokine production in mouse asthma model. *Cytokine*. 2013;61:237–244.

Lehrer SB, McClain S. Utility of animal models for predicting human allergenicity. *Regul Toxicol Pharmacol.* 2009;54:S46–51. doi: 10.1016/j.yrtph.2009.01.001.

Lehrer SB, Reese G, Malo JL, Lahoud C, Leong-Kee S, Goldberg B, et al. Corn allergens: IgE antibody reactivity and cross-reactivity with rice, soy, and peanut. *Int Arch Allergy Immunol.* 1999;118:298–299.

Leloup, V.M., Colonna, P., Ring, S.G., Roberts, K. & Wells, B. (1992). Microstructure of amylose gels. *Carbohydrate Polymers*, 18, 189–197.

Leung PS, Lee YS, Tang CY, Kung WY, Chuang YH, Chiang BL, Fung MC, Chu KH. Induction of shrimp tropomyosin-specific hypersensitivity in mice. *Int Arch Allergy Immunol.* 2008;147:305–314. doi: 10.1159/000144038.

Levin B.E., Hogan S., Sullivan A.C. Initiation and perpetuation of obesity and obesity resistance in rats. *Am. J. Physiol.* 1989;256:R766–R771.

Lewis D.F. 57 varieties: The human cytochromes P450. *Pharmacogenomics*. 2004;5:305–318.

Lezaun A, Igea JM, Quirce S, Cuevas M, Parra F, Alonso MD, Martin JA, Cano MS. Asthma and contact urticaria caused by rice in a housewife. *Allergy*. 1994;49:92–95. doi: 10.1111/j.1398-9995.1994.tb00806.x.



- Li C.H., Evans H.M. The isolation of pituitary growth hormone. *Science*. 1944;99:183–184.
- Li Ming, Li Yun-gao, GAO Hong-yan Technique Center of Shanghai Liangyou Group, Shanghai 200136, China; School of Food Science and Technology. Southern Yangtze University, Jiangsu Wuxi 214036, China): Research on Rice Protein (J): *Cereals & Oils*: 2006-08.
- Li P, Zang W, Li Y, Xu F, Wang J, Shi T. AtPID: the overall hierarchical functional protein interaction network interface and analytic platform for Arabidopsis. *Nucleic Acids Res*. 2010;39:D1130–1133.
- Li W., Li L., Li K., Lin J., Sun X., Tang K. Expression of biologically active human insulin-like growth factor 1 in *Arabidopsis thaliana* seeds via oleosin fusion technology. *Biotechnol. Appl. Biochem*. 2011;58:139–146.
- Li XJ, Yang MF, Chen H, Qu LQ, Chen F, Shen SH. Abscisic acid pretreatment enhances salt tolerance of rice seedlings: proteomic evidence. *Biochim Biophys Acta*. 2010;1804(4):929–40.
- Li XM, Schofield BH, Huang CK, Kleiner GI, Sampson HA. A murine model of IgE-mediated cow's milk hypersensitivity. *J Allergy Clin Immunol*. 1999;103:206–214. doi: 10.1016/S0091-6749(99)70492-6.
- Liebming E., Huttner S., Vavra U., Fischl R., Schoberer J., Grass J., Blaukopf C., Seifert G.J., Altmann F., Mach L., et al. Class I alpha-mannosidases are required for N-glycan processing and root development in *Arabidopsis thaliana*. *Plant. Cell*. 2009;21:3850–3867.
- Lienard D., Sourrouille C., Gomord V., Faye L. Pharming and transgenic plants. *Biotechnol. Annu. Rev*. 2007;13:115–147.
- Liener, I.E., 1994. Implication of anti-nutritional components in soybeans foods. *Crit. Rev. Food Sci. Nutr.*, 34: 31-67.
- Lii, C.Y., Tsai, M.L. & Tseng, K.H. (1996). Effect of amylose content on the rheological property of rice starch. *Cereal Chemistry*, 73, 415–420. Composition and functional properties of rice Z. Zhou et al. 865 \_ 2002 Blackwell Science Ltd *International Journal of Food Science and Technology* 2002, 37, 849–868.
- Lim, S.T., Lee, J.H., Shin, D.H. & Lim, H.S. (1999). Comparison of protein extraction solutions for rice starch isolation and effects of residual protein content on starch pasting properties. *Starch/Starke*, 51, 120–125.
- Lin C., Lin X., Hu L., Yang J., Zhou T., Long L., Xu C., Xing S., Qi B., Dong Y., et al. Dramatic genotypic difference in, and effect of genetic crossing on, tissue culture-induced mobility of retrotransposon Tos17 in rice. *Plant. Cell Rep*. 2012;31:2057–2063.
- Lin, W.M.: change S.M. and iii, C.Y (1979) studies on the changes of physiochemical properties of rice ageing storage. *Bull inst chem. Academic sin* 26 pg. 13-23.
- Lin SS, Martin R, Mongrand S, Vandenabeele S, Chen KC, Jang IC, Chua NH. RING1 E3 ligase localizes to plasma membrane lipid rafts to trigger FB1-induced programmed cell death in *Arabidopsis*. *Plant J*. 2008;56:550–561. doi: 10.1111/j.1365-313X.2008.03625.x.

Lin M, Hu B, Chen L, Sun P, Fan Y, Wu P, Chen X. Computational identification of potential molecular interactions in Arabidopsis. *Plant Physiol.* 2009;151:34–46. doi: 10.1104/pp.109.141317.

Lin M, Shen X, Chen X. PAIR: the predicted Arabidopsis interactome resource. *Nucleic Acids Res.* 2011;39:D1134–1140. doi: 10.1093/nar/gkq938.

Lin S, Chang M, Tsai Y, Lur H. Proteomic analysis of the expression of proteins related to rice quality during caryopsis development and the effect of high temperature on expression. *Proteomics.* 2005;5(8):2140–2156. doi: 10.1002/pmic.200401105.

Lindhout T., Iqbal U., Willis L.M., Reid A.N., Li J., Liu X., Moreno M., Wakarchuk W.W. Site-specific enzymatic polysialylation of therapeutic proteins using bacterial enzymes. *Proc. Natl. Acad. Sci. USA.* 2011;108:7397–7402.

Lisle, A.J., Martin, M. & Fitzgerald, M.A. (2000). Chalky and translucent rice grains differ in starch composition and structure and cooking properties. *Cereal Chemistry*, 77, 627–632.

Liu F., Zhao Y.Y., Zheng J.G. High-level expression of modified gene encoding human adiponectin in transgenic rice. *Biol. Res.* 2011;44:369–375.

Liu, H., Arntfield, S.D., Holley, R.A. & Aime, D.B. (1997). Amylose-lipid complex formation in acetylated pea starch-lipid systems. *Cereal Chemistry*, 74, 159–162.

Liu Y.K., Huang L.F., Ho S.L., Liao C.Y., Liu H.Y., Lai Y.H., Yu S.M., Lu C.A. Production of mouse granulocyte-macrophage colony-stimulating factor by gateway technology and transgenic rice cell culture. *Biotechnol. Bioeng.* 2012;109:1239–1247.

Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods.* 2001;25:402–408. doi: 10.1006/meth.2001.1262.

Lo WMY, Li-Chan ECY: Angiotensin I converting enzyme inhibitory peptides from in vitro pepsin-pancreatin digestion of soy protein. *J Agric Food Chem* 2005, 53:3369-3376.

Lopes CT, Franz M, Kazi F, Donaldson SL, Morris Q, Bader GD. Cytoscape Web: an interactive web-based network browser. *Bioinformatics.* 2010;26:2347–2348. doi: 10.1093/bioinformatics/btq430.

Lowry OH, Rosebrough NJ, Farr AL, Randall RJ: Protein measurement with the folin phenol reagent. *J Nutr Biochem* 1951, 193:265-275.

Lu D., Yang C., Liu Z. How hydrophobicity and the glycosylation site of glycans affect protein folding and stability: A molecular dynamics simulation. *J. Phys. Chem. B.* 2012;116:390–400.

Lu, S., Chen, L.N. & Lii, C.Y. (1997). Correlations between the fine structure, physicochemical properties, and retrogradation of amylopectins from Taiwan rice varieties. *Cereal Chemistry*, 74, 34–39.

- Lu T.J., Chen H.N., Wang H.J. Chemical constituents, dietary fiber, and  $\gamma$ -oryzanol in six commercial varieties of brown rice from Taiwan. *Cereal Chem.* 2011;88:463–466.
- Ludwig D.S., Majzoub J.A., Al-Zahrani A., Dallal G.E., Blanco I., Roberts S.B. High glycemic index foods, overeating, and obesity. *Pediatrics.* 1999;103 doi: 10.1542/peds.103.3.e26.
- Luh, B.S. & Liu, Y.K. (1991). Rice flours in baking. In: *Rice: Utilization* (edited by B.S. Luh). Pp. 9–33. New York: Van. Nostrand Reinhold.
- Lumdubwong, N. & Seib, P.A. (2000). Rice starch isolation by alkaline protease digestion of wet-milled rice flour. *Journal of Cereal Science*, 31, 63–74.
- Lund, E. K., J . M. Gee, J . C. Brown, Wood, P. J., and Johnson, I. T. 1989. Effect of oat gum on the physical properties of the gastrointestinal contents and on the uptake of D\_galactose and cholesterol by rat small intestine in vitro. *Br. J. Nutr.* 62:91-101..
- Lyon, Champagne, T; vineyard, B.T. Windham, W. R; Barton F.E. ii, Webb, B.O. McClung. A. N; Moldenhalelr K. A; Linscombe S; McKenzie, K. S. et al. (1999) effect of degree of milling, drying condition and moisture content on sensory texture of cooked rice cereal chem. Vol. 76 Pg. 56-62.
- M. Frei and K. Becker, 2003, Studies on the in vitro starch digestibility and glycemic index of six different indigenous rice cultivars from the Philippines, *Journal of Food Chemistry*, 83, 395-400.
- M. Yousaf, 1992 Study on some Physico-chemical characteristics affecting cooking and eating qualities of some Pakistani Rice Varieties, M.Sc. Thesis Department of Food Technology, University of Agriculture Faisalabad, Pakistan.
- Maki, M and Collin, P, Coeliac Disease, *Lancet* 349:1755-1759, 1997.
- Majamaa H, Laine S, Miettinen A. Eosinophil protein X and eosinophil cationic protein as indicators of intestinal inflammation in infants with atopic eczema and food allergy. *Clin Exp Allergy.* 1999;29:1502–1506.
- Malinow, M. R., P. McLaughwin, L. Papworth, H. K. Naito, and L. A. Lewis. 1976. Effect of bran and cholestyramine on plasma lipids in monkeys. *Am. J. Clin. Nutr.* 29:905-911.
- Mangala, S.L., Udayasankar, K. & Tharanathan, R.N. (1999). Resistant starch from processed cereals: the influence of amylopectin and non-carbohydrate constituents in its formation. *Food Chemistry*, 64, 391–396.
- Maningat, C.C. & Juliano, B.O. (1980). Starch lipids and their effect on rice starch properties. *Starch/Starke*, 32, 76–82.
- Mannheim, A. and M. Cheryan, 1990. Continuous hydrolysis of milk protein in a membrane reactor. *J. Food Sci.*, 55: 381-385.

Mano, Y., Kawaminami, K., Kojima, M., Ohnishi, M. & Ito, S. (1999). Comparative composition of brown rice lipids (lipid fractions) of Indica and Japonica rices. *Bioscience Biotechnology and Biochemistry*, 63, 619–626.

Marmagne A, Ferro M, Meinel T, Bruley C, Kuhn L, Garin J, Barbier-Brygoo H, Ephritikhine G. A high content in lipid-modified peripheral proteins and integral receptor kinases features in the Arabidopsis plasma membrane proteome. *Mol Cell Proteomics*. 2007;6:1980. doi: 10.1074/mcp.M700099-MCP200.

Marsh, MN, Gluten, Major Histocompatibility Complex and the Small Intestine, *Gastroenterology*, 102:330-354, 1992.

Marsh, MN, ed *Celiac Disease Methods and Protocols*, 2000.

Marshall, W.E. (1992). Effect of degree of milling of brown rice and particle size of milled rice on starch gelatinization. *Cereal Chemistry*, 69, 632–636.

Marshall, W.E., Normand, F.L. & Goynes, W.R. (1990). Effects of lipid and protein removal on starch gelatinization in whole grain milled rice. *Cereal Chemistry*, 67, 458–463.

Martin, C. & Smith, A.M. (1995). Starch biosynthesis. *Plant Cell*, 7, 971–985. Matsue, Y. & Ogata, T. (1998). Studies on palatability of rice grown in northern Kyushu – comparison of palatability between old and new types of rice cultivars. *Japanese Journal of Crop Science*, 67, 312–317.

Maslov S, Sneppen K. Specificity and stability in topology of protein networks. *Science*. 2002;296:910–913. doi: 10.1126/science.1065103.

Matthews LR, Vaglio P, Reboul J, Ge H, Davis BP, Garrels J, Vincent S, Vidal M. Identification of potential interaction networks using sequence-based searches for conserved protein-protein interactions or "interologs". *Genome Res*. 2001;11:2120–2126. doi: 10.1101/gr.205301.

Mattson, F.H., S. M. Grundy, and J. R. Crouse. 1982. Optimizing the effect of plant sterols on cholesterol absorption in man. *Am. J. Clin. Nutr.* 35:697-700.

Maurice, T.J., Slade, L., Sirett, R.R. & Page, C.M. (1985). Polysaccharide–water interactions—thermal behaviour of rice starch. In: *Influence of Water on Food Quality and Stability* (edited by D. Simatua & S.L. Multon). Dordrecht, The Netherlands: Martinus-Nijhoff.

McCarthy FM, Wang N, Magee GB, Nanduri B, Lawrence ML, Camon EB, Barrell DG, Hill DP, Dolan ME, Williams WP. et al. AgBase: a functional genomics resource for agriculture. *BMC Genomics*. 2006;7:229. doi: 10.1186/1471-2164-7-229.

McCormick A.A., Reddy S., Reinl S.J., Cameron T.I., Czerwinski D.K., Vojdani F., Hanley K.M., Garger S.J., White E.L., Novak J., et al. Plant-produced idiotype vaccines for the treatment of non-Hodgkin's lymphoma: Safety and immunogenicity in a phase I clinical study. *Proc. Natl. Acad. Sci. USA*. 2008;105:10131–10136.

- McCready, R.M. & Hassid, W.H. (1943). The separation and quantitative estimation of amylose and amylopectin in starch. *Journal of the American Chemical Society*, 65, 1154–1157.
- McDonald K.A., Hong L.M., Trombly D.M., Xie Q., Jackman A.P. Production of human alpha-1-antitrypsin from transgenic rice cell culture in a membrane bioreactor. *Biotechnol. Prog.* 2005;21:728–734.
- Mead JR, Irvine SA, Ramji DP: Lipoprotein lipase: structure, function, regulation, and role in disease. *J Mol Med* 2002, 80:753-769.
- Mehr SS, Kakakios AM, Kemp AS. Rice: a common and severe cause of food protein-induced enterocolitis syndrome. *Arch Dis Child.* 2009;94:220–223. doi: 10.1136/adc.2008.145144.
- Menon B, Peegel H, Menon K. Evidence for the association of luteinizing hormone receptor mRNA-binding protein with the translating ribosomes Michelet B, Boutry M. The Plasma Membrane H<sup>+</sup>-ATPase (a highly regulated enzyme with multiple physiological functions) *Plant Physiol.* 1995;108(1):1–6.
- Metcalf D. The granulocyte-macrophage colony stimulating factors. *Cell.* 1985;43:5–6.
- Meullenet J.F.C. Marks. B.P. Griffin. K. and Damels M. J (1999) effects of rough rice drying and storage condition on sensory profiles of cooked rice cereal chemistry vol. 76 pg. 483-486.
- Meullenet J.F. Marks Hankins J.A. Grillin V.K. and Daniles, (2000) sensory quality of cooked long grain rice as affected by rough rice moisture content storage temperature and storage duration cereal chemistry vol. 77 pg. 259-263.
- Meullenet, J.F., Griffin, V.K., Carson, K. et al. (2001). Rice external preference mapping for Asian consumers living in the United States. *Journal of Sensory Studies*, 16, 73–94.
- Michelet B, Boutry M. The Plasma Membrane H<sup>+</sup>-ATPase (a highly regulated enzyme with multiple physiological functions) *Plant Physiol.* 1995;108(1):1–6.
- Miles, M.J., Morris, V.J. & Ring, S.G. (1985). Gelation of amylose. *Carbohydrate Research*, 136, 257–269.
- Mizukami, H., Takeda, Y. & Hizukuri, S. (1999). The structure of the hot-water soluble components in the starch granules of new Japanese rice cultivars. *Carbohydrate Polymers*, 38, 329–335.
- Miller J.B., Pang E., Bramall L. Rice: A high or low glycemic index food? *Am. J. Clin. Nutr.* 1992;56:1034–1036.
- Miller A., Engel K.H. Content of  $\gamma$ -oryzanol and composition of steryl ferulates in brown rice (*Oryza sativa* L.) of European origin. *J. Agric. Food Chem.* 2006;54:8127–8133.
- Mingwei Min, Haoyang Cai, Zheng Wen, Zhirui Yang, Xiao Li. PlaPID: a database of protein-protein interactions in plants. *Bioinformatics and Biomedical Engineering (iCBBE) 2010 4th International Conference.* 2010;10:1109.

Mishra GR, Suresh M, Kumaran K, Kannabiran N, Suresh S, Bala P, Shivakumar K, Anuradha N, Reddy R, Raghavan TM. et al. Human protein reference database--2006 update. *Nucleic Acids Res.* 2006;34:D411–414. doi: 10.1093/nar/gkj141.

Mitschke D., Reichel A., Fricker G., Moenning U. Characterization of cytochrome P450 protein expression along the entire length of the intestine of male and female rats. *Drug Metab. Dispos.* 2008;36:1039–1045.

Mittler R, Rizhsky L. Transgene-induced lesion mimic. *Plant Mol Biol.* 2000;44:335–344. doi: 10.1023/A:1026544625898.

Miyata S., Akazawa T. Enzymic mechanism of starch breakdown in germinating rice seeds: 12. Biosynthesis of alpha-amylase in relation to protein glycosylation. *Plant. Physiol.* 1982;70:147–153.

Miyoshi, H., T. Okuda, Y. Oi, and H. Koishi. 1986. Effects of rice fiber on fecal weight, apparent digestibility of energy, nitrogen and fat, and degradation of neutral detergent fiber in young men. *J. Nutr. Sci. VitnIninol.*32:581\_589.

Mohamed T.K., Zhu K., Issoufou A., Fatmata T., Zhou H.M. Functionality, in vitro digestibility and physicochemical properties of two varieties of defatted foxtail millet protein concentrates. *Int. J. Mol. Sci.* 2009;10:5224–5238.

Momma, M., Saito, M., Chikuni, K. & Saio, K. (2000). Ultrastructure of prolamin accumulating protein bodies in endosperm of mutant rice for storage proteins –(Ultrastructure of protein bodies in mutant rice). *Journal of the Japanese Society for Food Science and Technology-Nippon Shokuhin Kagaku Kogaku Kaishi*, 47, 938–942.

Monteiro AG, Aoki MS, Evangelista AL, Alveno DA, Monteiro GA, Picarro Ida C, Ugrinowitsch C: Nonlinear periodization maximizes strength gains in split resistance training routines. *Journal of strength and conditioning research / National Strength & Conditioning Association* 2009, 23:1321-1326.

Monzón S, Lombardero M, Pérez-Camo I, Sáenz D, Lasanta J. Allergic rhinoconjunctivitis after ingestion of boiled rice. *J Investig Allergol Clin Immunol.* 2008;18:487–488.

Moore DR, Robinson MJ, Fry JL, Tang JE, Glover EI, Wilkinson SB, Prior T, Tarnopolsky MA, Phillips SM: Ingested protein dose response of muscle and albumin protein synthesis after resistance exercise in young men. *Am J Clin Nutr* 2009, 89:161-168.

Morita, T., and S. Kiriya, Mass Production Method for Rice Protein Isolate and Nutritional Evaluation, *J. Food Sci.* 56:1393–1396 (1993).

Morita T., Oh-hashii A., Kasaoka S., Ikai M., Kiriya S. Rice protein isolates produced by the two different methods lower serum cholesterol concentration in rats compared with casein. *J. Sci. Food Agric.* 1996;71:415–424.

Morita T., Oh-hashii A., Takei K., Ikai M., Kasaoka S., Kiriyama S. Cholesterol-lowering effects of soybean, potato and rice proteins depend on their low methionine contents in rats fed a cholesterol-free purified diet. *J. Nutr.* 1997;127:470–477.

Morita, Y. and C. Yoshida. 1968. Studies on gamma globulin of rice embryo. Part I. Isolation and purification of gamma globulin from rice embryo. *J. Biol. Chem.*, 32: 664-670.

Moritaka, S. Swada, K and Yasumatsu, K. (1972) studies on cereal vii storage determination of defatted Rice Eiyo to shokuryo vol. 25 pg. 16-20.

Moritaka, S. Yasumatus K. (1972) studies on cerea x. the effect of sulthydry 1 group on storage determination of milled rice. *Rice Eiyo to shokuryo* vol. 25 pg 59-62.

Morrison, W.R. (1978). Cereal lipids. In: *Advances in Cereal Science and Technology* (edited by Y. Pomeranz). Pp. 221–348. St Paul, Minnesota: American Association of Cereal Chemists. Morrison, W.R. (1988). Lipids in cereal starches: a review. *Journal of Cereal Science*, 8, 1–15.

Morrison, W.R. (1995). Starch lipids and how they relate to starch granule structure and functionality. *Cereal Chemistry*, 40, 437–446.

Morrison, W.R., Law, R.V. & Snape, C.E. (1993). Evidence for inclusion complexes of lipids with V-amylose in maize, rice and oat starches. *Journal of Cereal Science*, 18, 107–109.

Murray E.E., Lotzer J., Eberle M. Codon usage in plant genes. *Nucleic Acids Res.* 1989;17:477–498.

Nadimpalli R, Yalpani N, Johal GS, Simmons CR. Prohibitins, stomatins, and plant disease response genes compose a protein superfamily that controls cell proliferation, ion channel regulation, and death. *J Biol Chem.* 2000;275:29579–29586. doi: 10.1074/jbc.M002339200.

Nakajima-Adachi H, Ebihara A, Kikuchi A, Ishida T, Sasaki K, Hirano K, Watanabe H, Asai K, Takahashi Y, Kanamori Y. et al. Food antigen causes TH2-dependent enteropathy followed by tissue repair in T-cell receptor transgenic mice. *J Allergy Clin Immunol.* 2006;117:1125–1132. doi: 10.1016/j.jaci.2006.01.016.

Nakase M, Usui Y, Alvarez-Nakase AM, Adachi T, Urisu A, Nakamura R, Aoki N, Kitajima K, Matsuda T. Cereal allergens: rice-seed allergens with structural similarity to wheat and barley allergens. *Allergy.* 1998;53:55–57.

Nambu M, Shintaku N, Ohta S. Rice allergy. *Pediatrics.* 2006;117:2331–2332.

Natera SHA, Ford KL, Cassin AM, Patterson JH, Newbiggin EJ, Bacic A. Analysis of the *Oryza sativa* plasma membrane proteome using combined protein and peptide fractionation approaches in conjunction with mass spectrometry. *J Proteome Res.* 2008;7:1159–1187. doi: 10.1021/pr070255c.

Naveeda, K. and J. Prakash, 2004. Nutritional quality of microwave-cooked and pressure-cooked legumes. *Int. J. Food Sci. Nutr.*, 55: 441-448.

Neelam Gurnani et al; Chemistry & Allied Sciences – Emerging Future Technologies to Boost Up Lives. ISSN: 2250-3013 IOSR Journal of Pharmacy, 90-96.

Newman, R. K., A. A. Betschart, C. W. Newman, and P. J. Hofer. 1992. Effect of full-fat or defatted rice bran on serum cholesterol. 42:37-43. *Plant Foods Hun., Nutr.*

Newsholme EA, Blomstrand E: Branched-chain amino acids and central fatigue. *J Nutr* 2006, 136:274S-276S.

Ng A, Bursteinas B, Gao Q, Mollison E, Zvelebil M. Resources for integrative systems biology: from data through databases to networks and dynamic system models. *Brief Bioinform.* 2006;7:318–330. doi: 10.1093/bib/bbl036.

Nicolosi R. J., L. M. Ausman, and D. M. Hegsted. 1991. Rice bran oil lowers serum total and low density lipoprotein cholesterol and apo B levels in nonhuman primates. *Atherosclerosis* 88:133-142.

Ning, H., J. Qiao, Z. Liu, Z. Lin, G. Li, Q. Wang, S. Wang and Y. Ding. 2010. Distribution of proteins and amino acids in milled and brown rice as affected by nitrogen fertilization and genotype. *J. Cereal Sci.*, 52: 90-95.

Nishimura A., Aichi I., Matsuoka M. A protocol for *Agrobacterium*-mediated transformation in rice. *Nat. Protoc.* 2006;1:2796–2802.

Niu X, Narasimhan M, Salzman R, Bressan RA, Hasegawa PM. NaCl regulation of plasma membrane H<sup>+</sup>-ATPase gene expression in a glycophyte and a halophyte. *Plant Physiol.* 1993;103(3):713–718. doi: 10.1104/pp.103.3.713.

Nobuo S. Cytochrome P450 changes in rats with streptozocin-induced diabetes. *Int. J. Biochem.* 1994;26:1261–1268.

Nohzadeh Malakshah S, Habibi Rezaei M, Heidari M, Hosseini Salekdeh G. Proteomics reveals new salt responsive proteins associated with rice plasma membrane. *Biosci Biotechnol Biochem.* 2007;71:2144–2154. doi: 10.1271/bbb.70027.

Noomhorm, A., Kongseree, N. & Apintanapong, N. (1997). Effect of aging on the quality of glutinous rice crackers. *Cereal Chemistry*, 74, 12–15.

Normand, F.L. & Marshall, W.E. (1989). Differential scanning calorimetry of whole grain milled rice and milled rice flour. *Cereal Chemistry*, 66, 317–320.

Normand, F.L., Soignet, D.M., Hogan, J.T. & Deobald, H.J. (1966). Content of certain nutrients and amino acid patterns in high-protein rice flour. *Rice Journal*, 69, 13–18.



Norton LE, Layman DK: Leucine regulates translation initiation of protein synthesis in skeletal muscle after exercise. *Nutr* 2006, 136:533S-537S.

Norton L, Wilson GJ: Optimal protein intake to maximize muscle protein synthesis. *AgroFood industry hi-tech* 2009, 20:54-57.

Norton LE, Wilson GJ, Layman DK, Moulton CJ, Garlick PJ: Leucine content of dietary proteins is a determinant of postprandial skeletal muscle protein synthesis in adult rats. *Nutr Metab* 2012, 9:67.

Norton LE, Wilson GJ, Rupassar I, Garlick PJ, Layman DK: Leucine contents of isonitrogenous protein sources predict post prandial skeletal muscle protein synthesis in rats fed a complete meal. *FASEB* 2009, 22:227.224.

Nowak-Wegrzyn A, Muraro A. Food protein-induced enterocolitis syndrome. *Curr Opin Allergy Clin Immunol.* 2009;9:371–377. doi: 10.1097/ACI.0b013e32832d6315.

Nykiforuk C.L., Shen Y., Murray E.W., Boothe J.G., Busseuil D., Rheaume E., Tardif J.C., Reid A., Moloney M.M. Expression and recovery of biologically active recombinant apolipoprotein AI(Milano) from transgenic safflower (*Carthamus tinctorius*) seeds. *Plant. Biotechnol. J.* 2011;9:250–263.

Nykiforuk C.L., Boothe J.G., Murray E.W., Keon R.G., Goren H.J., Markley N.A., Moloney M.M. Transgenic expression and recovery of biologically active recombinant human insulin from *Arabidopsis thaliana* seeds. *Plant Biotechnol. J.* 2006;4:77–85.

O'Brien KP, Remm M, Sonnhammer EL. Inparanoid: a comprehensive database of eukaryotic orthologs. *Nucleic Acids Res.* 2005;33:D476–480.

Obembe O.O., Popoola J.O., Leelavathi S., Reddy S.V. Advances in plant molecular farming. *Biotechnol. Adv.* 2011;29:210–222.

Obenauer JC, Cantley LC, Yaffe MB. Scansite 2.0: Proteome-wide prediction of cell signaling interactions using short sequence motifs. *Nucleic Acids Res.* 2003;31(13):3635–3641. doi: 10.1093/nar/gkg584.

Ocker, H.D.; Boiling H: El-baya; AW (1976) effect of parboiling on some vitamins and mineral of rice thamine, riboflavin, calcium, magnesium manganese and phosphorus. *Riso* pg. 25, 79-82.

Ogawa, M., T. Kumamaru, H. Satoh, N. Iwata, T. Omura, Z. Kasai and K. Tanaka. 1987. Purification of protein body-I of rice seed and its polypeptide composition. *Plant Cell Physiol.*, 28: 1517-1527.

Ogawa Y., Sakurai N., Oikawa A., Kai K., Morishita Y., Mori K., Moriya K., Fujii F., Aoki K., Suzuki H., et al. High-throughput cryopreservation of plant cell cultures for functional genomics. *Plant. Cell Physiol.* 2012;53:943–952.

Ogawa, W., Kumamaru, T., Satoh, H. et al. (1989). Mutants of rice storage proteins. Isolation and characterization of protein bodies from rice mutants. *Theoretical and Applied Genetics*, 78, 306–309.

Ohtsubo, K., Kobayashi, A. & Shimizu, H. (1993). Quality evaluation of rice in Japan. *Japan Agricultural Research Quarterly*, 27, 95–101.

Ohtsubo K, Nakamura S: Cultivar identification of rice (*Oryza sativa* L.) by polymerase chain reaction method and its application to processed rice products. *J Agric Food Chem* 2007, 55:1501-1509.

Okaka J.C (2005); hardly storage and processing of plant food O.J academic publisher pg. 16-39.

Okuzaki A., Konagaya K., Nanasato Y., Tsuda M., Tabei Y. Estrogen-inducible GFP expression patterns in rice (*Oryza sativa* L.) *Plant. Cell Rep.* 2011;30:529–538.

Ooi HS, Schneider G, Chan YL, Lim TT, Eisenhaber B, Eisenhaber F. Databases of protein-protein interactions and complexes. *Methods Mol Biol.* 2010;609:145–159. doi: 10.1007/978-1-60327-241-4\_9.

Optimized Stabilized Rice Bran Extract for Metabolic Syndrome, Rice Science LLC, Singapore. (2009).

Orford, P.D., Ring, S.G., Carroll, V., Miles, M.J. & Morris, V.J. (1987). The effect of concentration and botanical source on the gelation and retrogradation of starch. *Journal of the Science of Food and Agriculture*, 39, 169–173.

Orhan F, Sekerel BE. A case of isolated rice allergy. *Allergy.* 2003;58:456–457. doi: 10.1034/j.1398-9995.2003.00132.x.

Ostlund G, Schmitt T, Forslund K, Kostler T, Messina DN, Roopra S, Frings O, Sonnhammer EL. InParanoid 7: new algorithms and tools for eukaryotic orthology analysis. *Nucleic Acids Res.* 2010;38:D196–203. doi: 10.1093/nar/gkp931.

Owens I.S., Ritter J.K., Yeatman M.T., Chen F. The novelUGT1 gene complex links bilirubin, xenobiotics, and therapeutic drug metabolism by encoding UDP-glucuronosyltransferase isozymes with a common carboxyl terminus. *J. Pharmacokinet Biopharm.* 1996;24:491–508.

P. R. Jennings, W. R. Coffman, and H. E. Kauffman, 1979, *Rice Improvement*, International Rice Research Institute, Los Banos, Philippines, 186p.

Paddon-Jones D, Sheffield-Moore M, Zhang XJ, Volpi E, Wolf SE, Aarsland A, Ferrando AA, Wolfe RR: Amino acid ingestion improves muscle protein synthesis in the young and elderly. *Am J Physiol Endocrinol Metab* 2004, 286:E321-E328.

Padhye, V. and D. Salunkhe. 1979. Extraction and characterization of rice proteins. *Cereal Chem.*, 56: 389-393.

Padhye, V.W. & Salunkhe, D.K. (1979). Extraction and characterisation of rice protein. *Cereal Chemistry*, 56, 389–393.

Pal, V., Pandey, J.P. & Sah, P.C. (1999). Effect of degree of polish on proximate composition of milled rice. *Journal of Food Science and Technology – Mysore*, 36, 160–162. 866 Composition and functional

properties of rice Z. Zhou et al. *International Journal of Food Science and Technology* 2002, 37, 849–868  
\_ 2002 Blackwell Science Ltd.

Pali-Scholl I, Yildirim AO, Ackermann U, Knauer T, Becker C, Garn H, Renz H, Jensen-Jarolim E, Fehrenbach H. Anti-acids lead to immunological and morphological changes in the intestine of BALB/c mice similar to human food allergy. *Exp Toxicol Pathol.* 2008;60:337–345. doi: 0.1016/j.etp.2008.03.004.

Palmiano, E.P., B.O. Juliano. 1972. Physicochemical properties of Niigata waxy rices. *Agr. Biol. Chem.* 36: 157-159.

Panlasigui L.N., Thompson L.U. Blood glucose lowering effects of brown rice in normal and diabetic subjects. *Int. J. Food Sci. Nutr.* 2006;57:151–158.

Park C.-I., Lee S.-J., Kang S.-H., Jung H.-S., Kim D.-I., Lim S.-M. Fed-batch cultivation of transgenic rice cells for the production of hCTLA4Ig using concentrated amino acids. *Proc. Biochem.* 2010;45:67–74.

Park HY, Kang IS, Han JS, Lee CH, An G, Moon YH. OsDEG10 encoding a small RNA-binding protein is involved in abiotic stress signaling. *Biochem Bioph Res Co.* 2009;380(3):597–602. doi: 10.1016/j.bbrc.2009.01.131.

Park, J.K., Kim, S.S. & Kim, K.O. (2001). Effect of milling ratio on sensory properties of cooked rice and on physicochemical properties of milled and cooked rice. *Cereal Chemistry*, 78, 151–156.

Pasiakos SM, McClung HL, McClung JP, Margolis LM, Andersen NE, Cloutier GJ, Pikosky MA, Rood JC, Fielding RA, Young AJ: Leucine-enriched essential amino acid supplementation during moderate steady state exercise enhances postexercise muscle protein synthesis. *Am J Clin Nutr* 2011, 94:809-818.

Pastorello EA, Farioli L, Pravettoni V, Ispano M, Scibola E, Trambaioli C, et al. The maize major allergen, which is responsible for food-induced allergic reactions, is a lipid transfer protein. *J Allergy Clin Immunol.* 2000;106:744–751.

Patel, R.M. (2006) Antioxidant Capacity of Stabilized Rice Bran. [www.RiceBranTech.com](http://www.RiceBranTech.com).

Patil A, Nakamura H. Filtering high-throughput protein-protein interaction data using a combination of genomic features. *BMC Bioinformatics.* 2005;6:100. doi: 10.1186/1471-2105-6-1000.

Pearce, K.N., and J.E. Kinsella, Emulsifying Properties of Proteins: Evaluation of a Turbidimetric Technique, *J. Agric. Food Chem.* 26:716–723 (1978).

Pellegrini M, Marcotte EM, Thompson MJ, Eisenberg D, Yeates TO. Assigning protein functions by comparative genome analysis: protein phylogenetic profiles. *Proc Natl Acad Sci USA.* 1999;96:4285–4288. doi: 10.1073/pnas.96.8.4285.

Pelkonen O., Turpeinen M., Hakkola J., Honkakoski P., Hukkanen J., Raunio H. Inhibition and induction of human cytochrome P450 enzymes: Current status. *Arch. Toxicol.* 2008;82:667–715.

Peng S, Huang J, Sheehy JE, Visperas RM. Rice yields decline with higher night temperature from global warming. *Proc Natl Acad Sci USA.* 2004;101(27):9971–9975. doi: 10.1073/pnas.0403720101.

Perdon A. A. Marks, B.P. Siebenmorgen T.J. and acid, Reid N.B. (1997) effect of rough rice storage conditions on the amylograph cooking properties of medium grain rice co. *Bengal cereal chemistry vol. 74 pg. 864-867.*

Perdon A. A. Siebenmorgen T.J. Buescher, R. W. and Gbur, E.E. (1999) starch retrogradation texture of cooked milled rice during storage *Journal food science vol. 64 pg. 828-832.*

Perdon, A., Siebenmorgen, T.J. & Mauromoustakos, A. (2000). Glassy state transition and rice drying: development of a brown rice state diagram. *Cereal Chemistry, 77, 708–713.*

Perlak F.J., Fuchs R.L., Dean D.A., McPherson S.L., Fischhoff D.A. Modification of the coding sequence enhances plant expression of insect control protein genes. *Proc. Natl. Acad. Sci. USA.* 1991;88:3324–3328.

Pharmaceutical Research: A Mouse Model to Evaluate the Impact of Species, Sex and Lipid Load on Lymphatic Drug Transport (</news/article/1332621/Pharmaceutical-Research-A-Mouse-Model-to-Evaluate-the-Impact-of-Species-Sex>).

Pharmaceutical Research: Protein Antigen Absorption to the DDA/TDB Liposomal Adjuvant: Effect on Protein Structure, Stability, and Liposome Physicochemical Characteristics ([news/article/1160915/Pharmaceutical-Research-protein-Antigen-Absorption-to-the-DDA-TDB-Liposomal-Adjuvant-Effect](/news/article/1160915/Pharmaceutical-Research-protein-Antigen-Absorption-to-the-DDA-TDB-Liposomal-Adjuvant-Effect)).

PHD Technologies LLC. Testing for USDA approval of Rice bran in standard of identity meat products.

Phillips SM: Dietary protein requirements and adaptive advantages in athletes. *Br J Nutr* 2012, 108(Suppl 2):S158-S167.

Pieterse CMJ, Leon-Reyes A, Van der Ent S, Van Wees SCM. Networking by small-molecule hormones in plant immunity. *Nat Chem Biol.* 2009;5:308–316. doi: 10.1038/nchembio.164.

Plewczynski D, Ginalski K. The interactome: predicting the protein-protein interactions in cells. *Cell Mol Biol Lett.* 2009;14:1–22. doi: 10.2478/s11658-008-0024-7.

Pomeranz, Y. and R.L. Oryl, 1982. Rice processing and utilization. In: *CRC Handbook of Processing and Utilization in Agriculture. Vol. II.* CRC Press, West Palm Beach, FL.

Porta C., Lomonossoff G.P. Viruses as vectors for the expression of foreign sequences in plants. *Biotechnol. Genet. Eng. Rev.* 2002;19:245–291.

- Prakash, J., 1996. Rice bran proteins: Properties and food uses. *Crit. Rev. Food Sci. Nutr.*, 36: 537-552.
- Pulchinelli A Jr, Costa AMM, de Carvalho CV, de Souza NCN, Haidar MA, Andriolo A, da Silva IDG: Positive association of the hepatic lipase gene polymorphism c.514C > T with estrogen replacement therapy response. *Lipids Health Dis* 2011, 10:197.
- Quirk A.V., Geisow M.J., Woodrow J.R., Burton S.J., Wood P.C., Sutton A.D., Johnson R.A., Dodsworth N. Production of recombinant human serum albumin from *Saccharomyces cerevisiae*. *Biotechnol. Appl. Biochem.* 1989;11:273–287.
- Qureshi, A. A., B. A. Bradlow, W. A. Salser, and L. D. Brace. 1997. Novel tocotrienol of rice bran modulate cardiovascular disease risk parameters of hypercholesterolemic humans. *J. Nutr. Biochem.* 8:290-298.
- Quresli, A. A., S. A. Sami, W. A. Salser, and F. A. Khan. 2002. Dose-dependent suppression of serum cholesterol by tocotrienol-rich fraction (TRF25) of rice bran in hypercholesterolemic humans. *Atherosclerosis* 161:199-207.
- Radauer C, Breiteneder H. Evolutionary biology of plant food allergens. *J Allergy Clin Immunol.* 2007;120:518–525.
- Radhika R.K. Ali S. Z252 Bhattacharya, K.R (1993) the fine structure of rice starch amylopectin and its relation of the texture of cooked rice carbohydrate, poly mp. 22, 267, -275.
- Raghuram, T. C., D. B. Rao, and C. Rukrini. 1989. Studies on hypolipidemic effects of dietary rice bran oil in human subjects. *Nutr. Rep. Int.* 39:889-895.
- Rajnara K., M. C. Prabhakar, and D. R. Krishna. 2001. Influence of rice bran oil on serum lipid peroxides and lipids in human subjects. *Indian J. Physiol Pharmacol.* 45:442-444.
- Ramesh, M., Ali, S.Z. & Bhattacharya, K.R. (1999a). Influence of parboiling on rice starch structure and cooked-rice texture. *Journal of Texture Studies*, 30, 337–344.
- Ramesh, M., Ali, S.Z. & Bhattacharya, K.R. (1999b). Structure of rice starch and its relation to cooked-rice texture. *Carbohydrate Polymers*, 38, 337–347.
- Ramesh, M., Mitchell, J.R., Jumel, K. & Harding, S.E. (1999c). Amylose content of rice starch. *Starch/Stärke*, 51, 311–313.
- Ramezanzadeh, F.M., R.M. Rao, P. Witoon, W.E. Marshall and M. Windhauser, 2000. Effects of microwave heat, packaging and storage temperature on fatty acid and proximate compositions in rice bran. *J. Agric. Food Chem.*, 48:464-467.
- Randall, J 61:114-116. . M., R. N. Sayre, W. G. Schultz, R. Y. Fong, A. P. Mossman, R. E. Tnbelhorn, and R. M. Saunders. 1985. Rice bran stabilization by extrusion cooking for extraction of edible oil. I. *Food Sci.* 50:361-364.

- Ranhotra, G. S., J. A. Gelroth, R. D. Reeves, M. K. Rudd, W. R. Durkee, J. D. Gardner. 1989. Short-term lipidemic responses in otherwise healthy hypercholesterolemic men consuming foods high in soluble fiber. *Cereal Client*. 66:94-97.
- Rasmussen BB, Phillips SM: Contractile and nutritional regulation of human muscle growth. *Exerc Sport Sci Rev* 2003, 31:127-131.
- Ravas E, Somera AL, Mongru DA, Oltvai ZN, Barabasi AL. Hierarchical organization of modularity in metabolic networks. *Science*. 2002;297:1551–1555. doi: 10.1126/science.1073374.
- Reddy, K.R., Ali, S.Z. & Bhattacharya, K.R. (1993). The fine structure of rice-starch amylopectin and its relation to the texture of cooked rice. *Carbohydrate Polymers*, 22, 267–275.
- Reeves PG, Nielsen FH, Fahey GC: AIN-93 purified diets for laboratory rodents: final report of the American Institute of Nutrition Ad Hoc Writing Committee on the reformulation of the AIN-76A rodent diet. *J Nutr* 1993, 123:1939-1951.
- Rehman, M.Y.A. and S. Mahmood, 1996. Anti-nutritional factors, metabolizable energy and chemical composition of rice bran. *MARDI Res. J.*, 24: 135-145.
- Remans T, Smeets K, Opdenakker K, Mathijsen D, Vangronsveld J, Cuypers A. Normalisation of real-time RT-PCR gene expression measurements in *Arabidopsis thaliana* exposed to increased metal concentrations. *Planta*. 2008;227:1343–1349. doi: 10.1007/s00425-008-0706-4.
- Ren J, Wen L, Gao X, Jin C, Xue Y, Yao X. CSS-Palm 2.0: an updated software for palmitoylation sites prediction. *Protein Eng Des Sel*. 2008;21:639–644. doi: 10.1093/protein/gzn039.
- Resurrección, A.P., Juliano, B.O. & Tanaka, Y. (1979). Nutrient content and distribution in milling fractions of rice grain. *Journal of the Science of Food and Agriculture*, 30, 475–481.
- Rhee SY, Beavis W, Berardini TZ, Chen G, Dixon D, Doyle A, Garcia-Hernandez M, Huala E, Lander G, Montoya M. et al. The *Arabidopsis* Information Resource (TAIR): a model organism database providing a centralized, curated gateway to *Arabidopsis* biology, research materials and community. *Nucleic Acids Res*. 2003;31:224–228. doi: 10.1093/nar/gkg076.
- Riego GD, Casano LM, Marti'n M, Sabater B. Multiple phosphorylation sites in the beta subunit of thylakoid ATP synthase. *Photosynth Res*. 2006;89(1):11–18. doi: 10.1007/s11120-006-9078-4.
- Riva, M., Schiraldi, A. & Piazza, L. (1994). Characterization of rice cooking – isothermal differential scanning calorimetry investigations. *Thermochimica Acta*, 246, 317–328.
- Rong, N. L., L. M. Ausman, and R. J. Nicolosi. 1997 Oryzanol decreases cholesterol absorption and aortic fatty streaks in hamsters. *Lipids*. 32:303-309.

- Rooke L, Byrne D, Salgueiro S. Marker gene expression driven by the maize ubiquitin promoter in transgenic wheat. *Ann Appl Biol.* 2000;136:167–172. doi: 10.1111/j.1744-7348.2000.tb00022.x.
- Roohinejad S., Mirhosseini H., Saari N., Mustafa S., Alias I., Shobirin M.H.A., Hamid A., Manap M.Y. Evaluation of GABA, crude protein and amino acid composition from different varieties of Malaysian's brown rice. *Aust. J. Crop Sci.* 2009;3:184–190.
- Rostoks N, Schmierer D, Kudrna D, Kleinhofs A. Barley putative hypersensitive induced reaction genes: genetic mapping, sequence analyses and differential expression in disease lesion mimic mutants. *Theor Appl Genet.* 2003;107:1094–1101. doi: 10.1007/s00122-003-1351-8.
- Rouanet, J-M, C. Laurent, and P. Besancon. 1993. Rice bran and wheat bran: selective effect on plasma and liver cholesterol in high-cholesterol fed rats. *Food Chem.* 47:67-71.
- Rual JF, Venkatesan K, Hao T, Hirozane-Kishikawa T, Dricot A, Li N, Berriz GF, Gibbons FD, Dreze M, Ayivi-Guedehoussou N. et al. Towards a proteome-scale map of the human protein-protein interaction network. *Nature.* 2005;437:1173–1178. doi: 10.1038/nature04209.
- Ruggiero P., Flati S., Di Cioccio V., Maurizi G., Macchia G., Facchin A., Anacardio R., Maras A., Lucarelli M., Boraschi D. Glycosylation enhances functional stability of the chemotactic cytokine CCL2. *Eur. Cytokine Netw.* 2003;14:91–96.
- Sahi C, Garwal M, Ingh A, Grover A. Molecular characterization of a novel isoform of rice (*Oryza sativa* L.) glycine rich-RNA binding protein and evidence for its involvement in high temperature stress response. *Plant Sci.* 2007;173:144–155. doi: 10.1016/j.plantsci.2007.04.010.
- Sakuma T., Honma R., Maguchi S., Tamaki H., Nemoto N. Different expression of hepatic and renal cytochrome P450s between the streptozotocin-induced diabetic mouse and rat. *Xenobiotica.* 2001;4:223–237.
- Saloga J, Renz H, Lack G, Bradley KL, Greenstein JL, Larsen G, Gelfand EW. Development and transfer of immediate cutaneous hypersensitivity in mice exposed to aerosolized antigen. *J Clin Invest.* 1993;91:133–140. doi: 10.1172/JCI116162.
- Salunkhe, D.K., J.K. Chavan, R.N. Adsule and S.S. Kadam, 1992. Rice. pp: 424-448. In: *World oilseeds: Chemistry, technology and utilization.* Van Nostrand Reinhold, New York.
- Salwinski L, Miller CS, Smith AJ, Pettit FK, Bowie JU, Eisenberg D. The Database of Interacting Proteins: 2004 update. *Nucleic Acids Res.* 2004;32:D449–451. doi: 10.1093/nar/gkh086.
- Sambrook J, Russell DW. *Molecular Cloning: a Laboratory Manual.* Cold Spring Harbor: Cold Spring Harbor Laboratory Press; 2001.
- Sanders, T. A. B., and S. Reddy. 1992. The influence of rice bran on plasma lipids and lipoproteins in human volunteers Eu,-. *I. Cl/n. Nut r.* 46:167-172.

Santamarina-Fojo S, González-Navarro H, Freeman L, Wagner E, Nong Z: Hepatic lipase, lipoprotein metabolism, and atherogenesis. *Arterioscler Thromb Vasc Biol* 2004, 24:1750-1754.

Sardana R.K., Alli Z., Dudani A., Tackaberry E., Panahi M., Narayanan M., Ganz P., Altosaar I. Biological activity of human granulocyte-macrophage colony stimulating factor is maintained in a fusion with seed glutelin peptide. *Transgenic Res.* 2002;11:521–531.

Sardana R., Dudani A.K., Tackaberry E., Alli Z., Porter S., Rowlandson K., Ganz P., Altosaar I. Biologically active human GM-CSF produced in the seeds of transgenic rice plants. *Transgenic Res.* 2007;16:713–721.

Saunders, R.M., 1990. Properties of rice bran as foodstuff. *Cereal Foods World*, 35: 632-635.

Savadkoobi, Sobhan; Hoogenkamp, Henk. Mechanical and gelling properties of comminuted sausage containing chicken MDM. Elsevier, *Journal of Food Engineering*. 117 (2013) 255-262.

Sawai, N., and Y. Morita, Studies on Rice Glutelin. II. Cross-Structure of Glutelin from Rice Endosperm, *Agr. Biol. Chem.(Tokyo)* 32:496–500 (1968).

Sayre, R.N., L. Earl, F.H. Kratzer and R.M. Saunders, 1987. Nutritional qualities of stabilized and raw rice bran for chicks. *Poult. Sci.*, 66: 493-499. Sayre, R. N., L. Earl, F. H. Kratzer, and R. M. Saunders. 1987 Nutritional qualities of stabilized and raw rice bran for chicks. *Poult Sci* 66:493-499.

Sayre, R. N., L. Earl, F. H. Kratzer, and R. M. Saunders. 1988. Effects of diets containing raw and extrusion-cooked rice bran on growth and efficiency of food utilization of broilers. *Brit. Poultry Sc.* 29:815-823.

Scafaro AP, Haynes PA, Atwell BJ. Physiological and molecular changes in *Oryza meridionalis* Ng., a heat-tolerant species of wild rice. *J Exp Bot.* 2010;61(1):191–202. doi: 10.1093/jxb/erp294.

Schagger, H., and G. von Jagow, Tricine-Sodium Dodecyl Sulfate- Polyacrylamide Gel Electrophoresis for the Separation of Proteins in the Range from 1 to 100 kDa, *Anal. Biochem.* 166: 368–379 (1987).

Schmider J., Greenblatt D.J., von Moltke L.L., Karsov D., Shader R.I. Inhibition of CYP2C9 by selective serotonin reuptake inhibitors in vitro: Studies of phenytoin p-hydroxylation. *Br. J. Clin. Pharmacol.* 1997;44:495–498.

Schroder K., Hertzog P.J., Ravasi T., Hume D.A. Interferon-gamma: An overview of signals, mechanisms and functions. *J. Leukoc. Biol.* 2004;75:163–189.

Seena, S., K.R. Sridhar, A.B. Arunb and C.C. Young, 2006. Effect of roasting and pressure-cooking on nutritional and protein quality of seeds of mangrove legume *Canavalia cathartica* from southwest coast of India. *J. Food Comp. Anal.*, 19: 284-293.

Seetharamaiah, G. S., and N. Chandrasekhara. 1989. Studies on hypocholesterolemic activity of rice bran oil. *Atherosclerosis* 78:219-223.



- Seetharamaiah, G. S., and N. Chandrasekhara. 1990. Effect of oryzanol on cholesterol absorption & biliary & fecal bile acids in rats. *Indian J. Med. Res.* [B] 92:471-475.
- Seetharamaiah, G. S., T. P. Krishnakantha, and N. Chandrasekhara. 1990. Influence of oryzanol on platelet aggregation in rats. *J. Nutr. Sci. Vitaminol.* 36:291-295.
- Sekhon, K.S., S.S. Dhillon, N. Singh and B. Singh, 1997. Functional suitability of commercially milled rice bran in India for use in different food products. *Plant Foods Hum. Nutr.*, 50: 127-140.
- Seow, C.C., Teo, C.H. & Nair, C.K.V. (1996). A DSC study of the effects of sugars on thermal properties of rice starch gels before and after aging. *Journal of Thermal Analysis*, 47, 1201–1212.
- Sesame and Rice Bran Oil Lowers Blood Pressure, Improves Cholesterol([/news/article/1172762/Sesame-and-Rice-Bran-Oil-Lowers-Blood-pressure-Improves-Cholesterol](#)).
- Sharif, K., M.S. Butt and N. Huma, 2005. Oil extraction from rice industrial waste and its effect on physicochemical characteristics of cookies. *Nutr. Food Sci.*, 35: 416-427.
- Sharma, H.R., G.S. Chauhan and K. Agrawal, 2004. Physico-chemical characteristics of rice bran processed by dry heating and extrusion cooking. *Int. J. Food Prop.*, 7: 603-614.
- Sharma, R. D., C. Rukmini. 1986. Rice bran oil and hypocholesterolemia in rats. *Lipids* 21:715-717.
- Sharma R. D., and C. Rukmini. 1987. Hypocholesterolemic activity of unsaponifiable matter of rice bran oil. *Indian J. Med. Res.* 85:278-281.
- She P, Horn CV, Reid T, Hutson SM, Cooney RN, Lynch CJ: Obesity-related elevations in plasma leucine are associated with alterations in enzymes involved in branched chain amino acid (BCAA) metabolism. *Am J Physiol Endocrinol Metab* 2007, 293:E1552-E1563.
- Shelness GS, Sellers JA: Very low density lipoprotein assembly and secretion. *Curr Opin Lipidol* 2001, 12:151-157.
- Shi, Y.-C. & Seib, P.A. (1992). The structure of four waxy starches related to gelatinization and retrogradation. *Carbohydrate Research*, 227, 131–145.
- Shibasaki M, Suzuki S, Nemoto H, Kuroume T. Allergenicity and lymphocyte-stimulating property of rice protein. *J Allergy Clin Immunol.* 1979;64:259–265. doi: 10.1016/0091-6749(79)90141-6.
- Shibuya, N; Iwasaki Y. Tanase, H. and Chikubu S. (1974) studies on deterioration of rice during storage. 1 change of brown rice and milled rice during storage . *Journal soc food science technology* vol. 21 pg. 597-603.
- Shibuya N. (1980) investigation on the deterioration of rice during storage. *Garin Journal (NaH grains authority manila)* vol. pg. 21-24.

- Shibuya, N'Iwasaki T and Chikubu. (1977) studies on deterioration of rice during storage iii on the changes of rice starch during storage of rice denpun kagaku vol. 24 pg. 55-58.
- Shih, F.F., 2003. An update on the processing of high-protein rice products. *Nahrung/Food*, 47: 420-424
- Steel, R.G.D., J.H. Torrie and D.A. Dickey, 1997. Principles and procedures of statistics: A biometrical approach, 3rd Ed. McGraw Hill Book Co., New York.
- Shih, F.F., E.T. Champagne, K. Daigle, and Z. Zarins, Use of Enzymes in the Processing of Protein Products from Rice Bran and Rice Flour, *Nahrung* 43:14–18 (1999).
- Shih, F. 2004. Rice proteins. In: *Rice Chemistry and Technology*. (Ed.): E.T. Champagne. AACC International, St. Paul, MN. pp. 143-157.
- Shih S.M., Doran P.M. Foreign protein production using plant cell and organ cultures: Advantages and limitations. *Biotechnol. Adv.* 2009;27:1036–1042.
- Shimojo N., Ishizaki T., Imaoka S., Funae Y., Fuji S., Okuda K. Changes in amounts of cytochrome P450 isozymes and levels of catalytic activities in hepatic and renal microsomes of rats with streptozocin-induced diabetes. *Biochem. Pharmacol.* 1993;46:621–627.
- Shin JE, Jeon YH, Yang HJ, Pyun BY. Comparison of clinical severity and laboratory results between atopic and non-atopic eczema in children. *Pediatr Allergy Respir Dis.* 2008;18:219–227.
- Shin Y.J., Hong S.Y., Kwon T.H., Jang Y.S., Yang M.S. High level of expression of recombinant human granulocyte-macrophage colony stimulating factor in transgenic rice cell suspension culture. *Biotechnol. Bioeng.* 2003;82:778–783.
- Shin Y.J., Chong Y.J., Yang M.S., Kwon T.H. Production of recombinant human granulocyte macrophage-colony stimulating factor in rice cell suspension culture with a human-like N-glycan structure. *Plant. Biotechnol. J.* 2011;9:1109–1119.
- Sievert, D. & Holm, J. (1993). Determination of amylose by differential scanning calorimetry. *Starch/Starke*, 45, 136–139.
- Singh, S., Dhaliwal, Y.S., Nagi, H.P.S. & Kalia, M. (1998). Quality characteristics of six rice varieties of Himachal Pradesh. *Journal of Food Science and Technology – Mysore*, 35, 74–78.
- Singh, V. (1998). Rice research finds no takers in the developed world. *The Economic Times*, 12 December, India.
- Singh, V., Okadome, H., Toyoshima, H., Isobe, S. & Ohtsubo, K. (2000). Thermal and physicochemical properties of rice grain, flour and starch. *Journal of Agricultural and Food Chemistry*, 48, 2639–2647.
- Sirko A., Vanek T., Gora-Sochacka A., Redkiewicz P. Recombinant cytokines from plants. *Int. J. Mol. Sci.* 2011;12:3536–3552.

Sivak, M.N. & Preiss, J. (1998). Starch: basic science to biotechnology. In: *Advances in Food and Nutrition Research* (edited by M.N. Sivak & J. Preiss). Pp. 18, 163–170. California: Academic Press.

Sivakumar G., Medina-Bolivar F., Lay J.O., Jr, Dolan M.C., Condori J., Grubbs S.K., Wright S.M., Baque M.A., Lee E.J., Paek K.Y. Bioprocess and bioreactor: Next generation technology for production of potential plant-based antidiabetic and antioxidant molecules. *Curr. Med. Chem.* 2011;18:79–90.

Skrabanek L, Saini HK, Bader GD, Enright AJ. Computational prediction of protein-protein interactions. *Mol Biotechnol.* 2008;38:1–17. doi: 10.1007/s12033-007-0069-2.

Smelko J.P., Wiltberger K.R., Hickman E.F., Morris B.J., Blackburn T.J., Ryll T. Performance of high intensity fed-batch mammalian cell cultures in disposable bioreactor systems. *Biotechnol. Prog.* 2011;27:1358–1364.

Smith, A.M., Denyer, K. & Martin, C. (1997). The synthesis of the starch granule. *Annual Review of Plant Physiology and Plant Molecular Biology*, 48, 67–87.

Smith JC, Fry AC, Weiss LW, Li Y, Kinzey SJ: The effects of high-intensity exercise on a 10-second sprint cycle test. *Journal of strength and conditioning research / National Strength & Conditioning Association* 2001, 15:344-348.

Sowbhagya, C.M., Ramesh, B.S. & Ali, S.Z. (1994). Hydration, swelling and solubility behaviour of rice in relation to other physicochemical properties. *Journal of the Science of Food and Agriculture*, 64, 1–7.

Spady, D. K., J. M. Dietsch y. 1985. Rates of cholesterol synthesis and low-density lipoprotein uptake in the adrenal glands of the rat, hamster and rabbit in vivo. *Biochim. Biophys. Acta* 836:167-175.

Sparrow P.A., Irwin J.A., Dale P.J., Twyman R.M., Ma J.K. Pharma-planta: Road testing the developing regulatory guidelines for plant-made pharmaceuticals. *Transgenic Res.* 2007;16:147–161.

Srinivasan K., Viswanad B., Asrat L., Kaul C.L., Ramarao P. Combination of high-fat diet-fed and low-dose streptozotocin-treated rat: A model for type 2 diabetes and pharmacological screening. *Pharmacol. Res.* 2005;52:313–320.

Sriraman R., Bardor M., Sack M., Vaquero C., Faye L., Fischer R., Finnern R., Lerouge P. Recombinant anti-HCG antibodies retained in the endoplasmic reticulum of transformed plants lack core-xylose and core-alpha(1,3)-fucose residues. *Plant. Biotechnol. J.* 2004;2:279–287.

Stark C, Breitkreutz BJ, Reguly T, Boucher L, Breitkreutz A, Tyers M. BioGRID: a general repository for interaction datasets. *Nucleic Acids Res.* 2006;34:D535–539. doi: 10.1093/nar/gkj109.

Steinhart, H., T. Rathjen. 2003. Dependence of tocopherol stability on different cooking procedures of food. *Int. J. Vitam. Nutr. Res.* 73:141-151.

Stelzl U, Worm U, Lalowski M, Haenig C, Brembeck FH, Goehler H, Stroedicke M, Zenkner M, Schoenherr A, Koeppen S. et al. A human protein-protein interaction network: a resource for annotating the proteome. *Cell*. 2005;122:957–968. doi: 10.1016/j.cell.2005.08.029.

Stoger E., Vaquero C., Torres E., Sack M., Nicholson L., Drossard J., Williams S., Keen D., Perrin Y., Christou P., et al. Cereal crops as viable production and storage systems for pharmaceutical ScFv antibodies. *Plant. Mol. Biol.* 2000;42:583–590.

Strid J, Thomson M, Hourihane J, Kimber I, Strobel S. A novel model of sensitization and oral tolerance to peanut protein. *Immunology*. 2004;113:293–303. doi: 10.1111/j.1365-2567.2004.01989.x.

Stumvoll M., Goldstein B.J., van Haeften T.W. Type 2 diabetes: Principles of pathogenesis and therapy. *Lancet*. 2005;365:1333–1346.

Sturgess, RP et al, Cereal Chemistry, Molecular Biology and Toxicity in Coeliac Disease, *Gut* 32:1055-1060, 1991.

Sturgess, RP et al, Wheat Peptide Challenge in Coeliac Disease, *Lancet*, 343:759-761, 1994.

Su C.F., Kuo I.C., Chen P.W., Huang C.H., Seow S.V., Chua K.Y., Yu S.M. Characterization of an immunomodulatory Der p 2-FIP-fve fusion protein produced in transformed rice suspension cell culture. *Transgenic Res*. 2012;21:177–192.

Su W.W., Arias R. Continuous plant cell perfusion culture: Bioreactor characterization and secreted enzyme production. *J. Biosci. Bioeng.* 2003;95:13–20.

Sugano, M., N. Ishiwaki, and K. Nakashima. 1984. Dietary protein-dependent modification of serum cholesterol level in rats. *Ann. Nutr. Metb.* 28:192-199. S

ugano M., Goto S., Yamada Y., Yoshida K., Hashimoto Y., Matsuo T., Kimoto M. Cholesterol-lowering activity of various undigested fractions of soybean protein in rats. *J. Nutr.* 1990;120:977–985.

Sugimoto, T., K. Tanaka and Z. Kasai. 1986. Improved extraction of rice prolamin. *Agric. Biol. Chem.*, 50: 2409-2411.

Sugio S., Kashima A., Mochizuki S., Noda M., Kobayashi K. Crystal structure of human serum albumin at 2.5 Å resolution. *Protein Eng.* 1999;12:439–446.

Sun Q., Donna S., Rob M.V.D., Michelle D.H., Vasanti S.M., Walter C.W., Frank B.H. White rice, brown rice, and risk of type 2 diabetes in US men and women. *Arch. Intern. Med.* 2010;170:961–969.

Sun Q.Y., Ding L.W., Lomonossoff G.P., Sun Y.B., Luo M., Li C.Q., Jiang L., Xu Z.F. Improved expression and purification of recombinant human serum albumin from transgenic tobacco suspension culture. *J. Biotechnol.* 2011;155:164–172.

- Sunitha, L., R. Manorama, and C. Rukmini. 1997 Lipid profile of rats fed blends of rice bran oil in combination with sunflower and safflower oil. *Plant Foods for Human Nutr.* 51:219-230.
- Suzanne D, Donald KL: Protein metabolic roles in treatment of obesity. *Curr Opin Clin Nutr Metab Care* 2010, 13:403-407.
- Suzuki, M. 1982. Repressive effect of dietary fiber fractions in unpolished rice on the increase in cholesterol and triglyceride. I. *Nutr. Food (Japan)* 35:155-160.
- Suzuki K., Kaminuma O., Yang L., Takai T., Mori A., Umezu-Goto M., Ohtomo T., Ohmachi Y., Noda Y., Hirose S., et al. Prevention of allergic asthma by vaccination with transgenic rice seed expressing mite allergen: Induction of allergen-specific oral tolerance without bystander suppression. *Plant Biotechnol. J.* 2011;9:982–990.
- Suzuki, S., and S. Oshima. 1970. Influence of blending of edible fats and oils on human serum cholesterol level (Part 1). Blending of rice bran oil and safflower oil. I. *Nutr. (Japan)*. 28:3-6.
- Suzuki, S., and S. Oshima. 1970. Influence of blending oils on human serum cholesterol (Part 2). Rice bran oil, safflower oil, sunflower oil. 28:194-198. *J. Nutr. (Japan)*.
- Swindell WR, Huebner M, Weber AP. Transcriptional profiling of Arabidopsis heat shock proteins and transcription factors reveals extensive overlap between heat and non-heat stress response pathways. *BMC Genomics*. 2007;8:125. doi: 10.1186/1471-2164-8-125.
- Tachibana N, Matsumoto I, Fukui K, Arai S, Kato H, Abe K, Takamatsu K: Intake of soy protein isolate alters hepatic gene expression in rats. *J Agric Food Chem* 2005, 53:4253-4257.
- Taira, H., Nakagahra, M. & Nagamine, T. (1988). Fatty acid composition of indica, sinica, japonica, and japonica groups of nonglutinous brown rice. *Journal of Agricultural and Food Chemistry*, 36, 45–47.
- Takagi H., Hiroi T., Yang L., Tada Y., Yuki Y., Takamura K., Ishimitsu R., Kawauchi H., Kiyono H., Takaiwa F. A rice-based edible vaccine expressing multiple T cell epitopes induces oral tolerance for inhibition of Th2-mediated IgE responses. *Proc. Natl. Acad. Sci. USA*. 2005;102:17525–17530.
- Takaiwa F. Allergen-specific immunotherapy with plant-based oral vaccines. *Immunotherapy*. 2009;1:517–519.
- Takeda, Y., Hizukuri, S. & Juliano, B.O. (1986). Purification and structure of amylose from rice starch. *Carbohydrate Research*, 148, 299–308.
- Takeda, Y., Hizukuri, S. & Juliano, B.O. (1989). Structures and amount of branched molecules in rice amylose. *Carbohydrate Research*, 186, 163–168.
- Takemasa, M. and S. Hijikuro, 1991. Effect of autoclaving of defatted rice bran on the phosphorus utilization for chicks. *Jap. Poult. Sci.*, 28: 284-293.

Tako, M. & Hizukuri, S. (1999). Gelatinization mechanism of rice starch. *Journal of Carbohydrate Chemistry*, 18, 573–584.

Tan, N.H., Z.H.A. Rahim, H.T. Khor and K.C. Wong, 1983. Winged bean (*Psophocarpustetragonolobus*) tannin level, phytate content and haemagglutinin activity. *J. Agric. Food Chem.*, 31: 591-599. Tan, Y.F., Li, J.X., Yu, S.B., Xing, Y.Z., Xu, C.G. & Zhang, Q. (1999). The three important traits for cooking and eating quality of rice grains are controlled by a single locus in an elite rice hybrid, Shanyou 63. *Theoretical and Applied Genetics*, 99, 642–648.

Tan, Y.F., Li, J.X., Yu, S.B., Xing, Y.Z., Xu, C.G. & Zhang, Q. (1999). The three important traits for cooking and eating quality of rice grains are controlled by a single locus in an elite rice hybrid, Shanyou 63. *Theoretical and Applied Genetics*, 99, 642–648.

Tanaka, K., Sugimoto, T., Ogawa, M. & Kasai, Z. (1980). Isolation and characterization of two types of protein bodies in the rice endosperm. *Agricultural and Biological Chemistry*, 44, 1633–1639.

Tang, S., N.S. Hettiarachchy and T.H. Shellhammer, 2002. Protein extraction from heat-stabilized defatted rice bran. I. Physical processing and enzyme treatments. *J. Agric. Food Chem.*, 50: 7444-7448.

Tang, S., N.S. Hettiarachchy, S. Eswaranandam and P. Crandall, 2003. Protein extraction from heatstabilized defatted rice bran: II. The role of amylase, celluclast and viscozyme. *J. Food Sci.*, 68: 471-475.

Tangendjaja, B., K.A. Buckle and M. Wootton, 2006. Dephosphorylation of phytic acid in rice bran. *J. Food Sci.*, 46: 1021-1024.

Tanaka, K., T. Sugimoto, M. Ogawa and Z. Kasai. 1980. Isolation and characterization of two types of protein bodies in the rice endosperm. *Agri. Biol. Chem.*, 44: 1633-1639.

Tang JE, Moore DR, Kujbida GW, Tarnopolsky MA, Phillips SM: Ingestion of whey hydrolysate, casein, or soy protein isolate: effects on mixed muscle protein synthesis at rest and following resistance exercise in young men. *J Appl Physiol* 2009, 107:987-992.

Tashiro, M. and S. Ikegami, 1996. Changes in activity, antigenicity and molecular size of rice bran trypsin inhibitor by in-vitro digestion. *Nutr. Sci. Vit.*, 42: 367-376.

Tacson, E., B. Esmama, L. Lontok and B. Juliano. 1971. Studies on the extraction and composition of rice endosperm glutelin and prolamin. *Cereal Chem.*, 48: 168-181.

Tester, R.F. & Morrison, W.R. (1990a). Swelling and gelatinization of cereal starches. I. Effects of amylopectin, amylose, and lipids. *Cereal Chemistry*, 67, 551–557.

Tester, R.F. & Morrison, W.R. (1990b). Swelling and gelatinization of cereal starches. II. Waxy rice starches. *Cereal Chemistry*, 67, 558–563.

- Thomma BPHJ, Eggermont K, Penninckx IAMA, Mauch-Mani B, Vogelsang R, Cammue B, Broekaert WF. Separate jasmonate-dependent and salicylate-dependent defense-response pathways in Arabidopsis are essential for resistance to distinct microbial pathogens. *Proc Natl Acad Sci USA*. 1998;95:15107–15111. doi: 10.1073/pnas.95.25.15107.
- Thomas JB, Robert E, Sharwood RG, Silva LM, David BS. The RNA-binding proteins CSP41a and CSP41b may regulate transcription and translation of chloroplast-encoded RNAs in Arabidopsis. *Plant Mol Biol*. 2009;69(5):541–552. doi: 10.1007/s11103-008-9436-z.
- Tipton KD, Ferrando AA, Phillips SM, Doyle D Jr, Wolfe RR: Postexercise net protein synthesis in human muscle from orally administered amino acids. *Am J Physiol* 1999, 276:E628-E634.
- Topping D. L., R. J. Illman, P. D. Roach, R. P. Trimble, A. Kambouris, and P. J. Nestel. 1990. Modulation of the hypolipidemic effect of fish oils by dietary fiber in rats: studies with rice and wheat bran. *J. Nutr.* 120:325-330.
- Torres E., Vaquero C., Nicholson L., Sack M., Stoger E., Drossard J., Christou P., Fischer R., Perrin Y. Rice cell culture as an alternative production system for functional diagnostic and therapeutic antibodies. *Transgenic Res*. 1999;8:441–449.
- Tran TU, Suzuki K, Okadome H, Ikezaki H, Homma S, Ohtsubo K: Detection of changes in taste of japonica and indica brown and milled rice (*Oryza sativa* L.) during storage using physicochemical analyses and a taste sensing system. *J Agric Food Chem* 2005, 53:1108-1118.
- Travis J., Salvesen G.S. Human plasma proteinase inhibitors. *Annu. Rev. Biochem.* 1983;52:655–709.
- Trexler M.M., McDonald K.A., Jackman A.P. Bioreactor production of human alpha(1)-antitrypsin using metabolically regulated plant cell cultures. *Biotechnol. Prog.* 2002;18:501–508.
- Trier, JS, Celiac Sprue, *New England Journal of Medicine*, 325:1709-1719, 1991.
- Triguero A., Cabrera G., Rodriguez M., Soto J., Zamora Y., Perez M., Wormald M.R., Cremata J.A. Differential N-glycosylation of a monoclonal antibody expressed in tobacco leaves with and without endoplasmic reticulum retention signal apparently induces similar in vivo stability in mice. *Plant. Biotechnol. J.* 2011;9:1120–1130.
- Trowell, H. C. 1975. *Refined Foods and Disease*, Burkitt and Trowell, Eds., Academic Press, London, 195-226.
- Tsai, C. E., H. Ting, L. J. Wang, and T. Lin. 1992. The effect of rice bran on blood lipids in man. *J. Chinese Agric. Chen. Soc.* 30:484-495.
- Turner A: The science and practice of periodization: a brief review. *Strength and Conditioning Journal* 2011, 33:34-46.

Untersmayr E, Jensen-Jarolim E. The role of protein digestibility and antacids on food allergy outcomes. *J Allergy Clin Immunol.* 2008;121:1301–1308. doi: 10.1016/j.jaci.2008.04.025. quiz 1309-1310.

Uetz P, Giot L, Cagney G, Mansfield TA, Judson RS, Knight JR, Lockshon D, Narayan V, Srinivasan M, Pochart P. et al. A comprehensive analysis of protein-protein interactions in *Saccharomyces cerevisiae*. *Nature.* 2000;403:623–627. doi: 10.1038/35001009.

Untersmayr E, Jensen-Jarolim E. The role of protein digestibility and antacids on food allergy outcomes. *J Allergy Clin Immunol.* 2008;121:1301–1308.

Urbano, G., M. Lopez-Jurado, J. Hernandez, M. Fernandez, M.C. Moreu, J. Frias, C.D. Pollan, M. Prodanov and C. Vidal-Valverde, 1995. Nutritional assessment of raw, heated and germinated lentils. *J. Agric. Food Chem.*, 43: 1871-1877.

USA Rice Federation, 2002, The natural history of rice, *Food and Cultural Encyclopedia*, 1-4.

Usui Y, Nakase M, Hotta H, Urisu A, Aoki N, Kitajima K, Matsuda T. A 33-kDa allergen from rice (*Oryza sativa* L. Japonica). cDNA cloning, expression, and identification as a novel glyoxalase I. *J Biol Chem.* 2001;276:11376–11381. doi: 10.1074/jbc.M010337200.

Vahouny G.V., Chalcarz W., Satchithanandam S., Adamson I., Klurfeld D.M., Kritchevsky D. Effect of soy protein and casein intake on intestinal absorption and lymphatic transport of cholesterol and oleic acid. *Am. J. Clin. Nutr.* 1984;40:1156–1164.

Vahouny G.V., Adamson I., Chalcarz W., Satchithanandam S., Muesing R., Klurfeld D.M., Tepper S.A., Sanghvi A., Kritchevsky D. Effects of casein and soy protein on hepatic and serum lipids and lipoprotein lipid distributions in the rat. *Atherosclerosis.* 1985;56:127–137.

Vailleau F, Daniel X, Tronchet M, Montillet JL, Triantaphylidès C, Roby D. A R2R3-MYB gene, AtMYB30, acts as a positive regulator of the hypersensitive cell death program in plants in response to pathogen attack. *Proc Natl Acad Sci USA.* 2002;99:10179–10184. doi: 10.1073/pnas.152047199.

Valeur J, Lappalainen J, Rita H, Lin AH, Kovanen PT, Berstad A, Eklund KK, Vaali K. Food allergy alters jejunal circular muscle contractility and induces local inflammatory cytokine expression in a mouse model. *BMC Gastroenterol.* 2009;9:33. doi: 10.1186/1471-230X-9-33.

Van Doorn WG, Woltering EJ. Many ways to exit? Cell death categories in plants. *Trends Plant Sci.* 2005;10:117–122. doi: 10.1016/j.tplants.2005.08.003.

Vasanathan, Y. & Hoover, R. (1992). Effect of defatting on starch structures and physicochemical properties. *Food Chemistry*, 45, 337–341.

Velasquez MT, Bhatena SJ: Role of dietary soy protein in obesity. *Int J Med Sci* 2007, 4:72-82. Villareal, C.P., Delacruz, N.M. & Juliano, B.O. (1994). Rice amylose analysis by near-infrared transmittance spectroscopy. *Cereal Chemistry*, 71, 292–296.



- Villareal, C.P., Hizukuri, S. & Juliano, B.O. (1997). Amylopectin staling of cooked milled rices and properties of amylopectin and amylose. *Cereal Chemistry*, 74, 163–167.
- Villareal, C.P. & Juliano, B.O. (1986). Waxy gene factor and residual protein of rice starch granules. *Starch/Starke*, 38,118–122.
- Villareal, C.P. & Juliano, B.O. (1989). Comparative levels of waxy gene product of endospermstarch granules of different rice ecotypes. *Starch/Starke*, 41, 369–373.
- Villareal, C.P., Juliano, B.O. & Hizukuri, S. (1993). Varietal differences in amylopectin staling of cooked waxy milled rices. *Cereal Chemistry*, 70, 753–758.
- Villareal, R.M. resurrection A.P. Suzuki, L.B. and Juliano, B.O. (1976) changes in physiochemical properties o rice during storage starch/starke vol. 28 pg. 88-94.
- Vinje NE, Larsen S, Lovik M. A mouse model of lupin allergy. *Clin Exp Allergy*. 2009;39:1255–1266. doi: 10.1111/j.1365-2222.2009.03269.x.
- Vitoria JC, Camarero C, Sojo A, Ruiz A, Rodriguez-Soriano J. Enteropathy related to fish, rice, and chicken. *Arch Dis Child*. 1982;57:44–48.
- Voinnet O., Rivas S., Mestre P., Baulcombe D. An enhanced transient expression system in plants based on suppression of gene silencing by the p19 protein of tomato bushy stunt virus. *Plant. J*. 2003;33:949–956.
- Wahlqvist ML, Hodgson JM, Ng FM, Hsu-Hage BHH, Strauss BJJ: The role of nutrition in abdominal obesity. *Nutr Res* 1999, 19:85-101.
- Wanasundara, P.K.J.P.D., and F. Shahidi, Removal of Flaxseed Mucilage by Chemical and Enzymatic Treatments, *Food Chem*. 59:47–55 (1997).
- Wang, M., N.S. Hettiarachchy, M. Qi, W. Burks and T. Siebenmorgen, 1999. Preparation and functional properties of rice bran protein isolate. *J. Agric. Food Chem.*, 47: 411-416. Wang YM, Zhang B, Xue Y, Li ZJ,
- Wang JF, Xue CH, Yanagita T: The mechanism of dietary cholesterol effects on lipids metabolism in rats. *Lipids Health Dis* 2010, 9:4.
- Wang H, Peng DQ: New insights into the mechanism of low high-density lipoprotein cholesterol in obesity. *Lipids Health Dis* 2011, 10:176.
- Wang H, Eckel RH: Lipoprotein lipase: from gene to obesity. *Am J Physiol Endocrinol Metab* 2009, 297:E271-E288.
- Wang YM, Zhang B, Xue Y, Li ZJ, Wang JF, Xue CH, Yanagita T. The mechanism of dietary cholesterol effects on lipids metabolism in rats. *Lipids Health Dis*. 2010;9:4. doi: 10.1186/1476-511X-9-4.

Wang H, Peng DQ. New insights into the mechanism of low high-density lipoprotein cholesterol in obesity. *Lipids Health Dis.* 2011;10:176. doi: 10.1186/1476-511X-10-176.

Wang Mei Shen hui Wuxi University of Light Industry Food School Wuxi. 214036, Bioactive Peptides in of Enzymatic Hydrolysate of Food Protein (J): *Amino Acids and Biotic Resources*: 1997-01.

Wang Zhang-cun. Shen Ru-ing. YAO Hui-yuan (School of Food Science & Technology, Southern Yangtze University, 214036 Wuxi, China). Extraction and Exploitation of Rice Protein (J), *Journal of Cereals & Oils*: 2004-01.

Wang Yalin, Zhong Fangxu, Tao Xingwu: Jiang Xianjun (Department of Biotechnology and Chemical Engineering. Wuhan Polytechnic University, Wuhan 430070, China): Study on the extraction of protein from rice dregs in alkalinous condition (J): *Journal of Wuhan Institute of Chemical Technology*.

Wang Z., Hall S.D., Maya J.F., Li L., Asghar A., Gorski J.C. Diabetes mellitus increases the in vivo activity of cytochrome P450 2E1 in humans. *Br. J. Clin. Pharmacol.* 2003;55:77–85.

Wang Y., Deng H., Zhang X., Xiao H., Jiang Y., Song Y., Fang L., Xiao S., Zhen Y., Chen H. Generation and immunogenicity of Japanese encephalitis virus envelope protein expressed in transgenic rice. *Biochem. Biophys. Res. Commun.* 2009;380:292–297.

Waskiewicz AJ, Johnson JC, Penn B, Mahalingam M, Kimball SR, Cooper JA. Phosphorylation of the cap-binding protein eukaryotic translation initiation factor 4E by protein kinase Mnk1 in vivo. *Mol Cellular Bio.* 1999;19(3):1871–1880.

Watts DJ, Strogatz SH. Collective dynamics of 'small-world' networks. *Nature.* 1998;393:440–442. doi: 10.1038/30918.

Weber, K. and M. Osborn. 1969. The reliability of molecular weight determinations by dodecyl sulfate-polyacrylamide gel electrophoresis. *J. Biol. Chem.*, 244: 4406-4412.

White, P.J., Abbas, I.R. & Johnson, L.A. (1989). Freezethaw stability and refrigerated-storage retrogradation of starches. *Starch/Stärke*, 41, 176–180.

Wichadakul D, McDermott J, Samudrala R. Prediction and integration of regulatory and protein-protein interactions. *Methods Mol Biol.* 2009;541:101–143. doi: 10.1007/978-1-59745-243-4\_6. *Am J Clin Nutr* 2007, 85:1031-1040.

Wilkinson SB, Tarnopolsky MA, Macdonald MJ, Macdonald JR, Armstrong D, Phillips SM: Consumption of fluid skim milk promotes greater muscle protein accretion after resistance exercise than does consumption of an isonitrogenous and isoenergetic soy-protein beverage.

Willis RC, Hogue CW. Searching, viewing, and visualizing data in the Biomolecular Interaction Network Database (BIND) *Curr Protoc Bioinformatics.* 2006;Chapter 8 Unit 8 9.

- Wiseman, M.O. and R.L. Price, 1987. Characterization of protein concentrates of jojoba (*Simonsia chinensis*) meal. *Cereal Chem.*, 64: 91-93.
- Wu H, Su Z, Mao F, Olman V, Xu Y. Prediction of functional modules based on comparative genome analysis and Gene Ontology application. *Nucleic Acids Res.* 2005;33:2822–2837. doi: 10.1093/nar/gki573.
- Wu KG, Li TH, Chen CJ, Cheng HI, Wang TY. Correlations of serum Interleukin-16, total IgE, eosinophil cationic protein and total eosinophil counts with disease activity in children with atopic dermatitis. *Int J Immunopathol Pharmacol.* 2011;24:15–23.
- Wu X, Zhu L, Guo J, Zhang DY, Lin K. Prediction of yeast protein-protein interaction network: insights from the Gene Ontology and annotations. *Nucleic Acids Res.* 2006;34:2137–2150. doi: 10.1093/nar/gkl219.
- Wu X, Zhu L, Guo J, Fu C, Zhou H, Dong D, Li Z, Zhang DY, Lin K. SPIDer: *Saccharomyces* protein-protein interaction database. *BMC Bioinformatics.* 2006;7(Suppl 5):S16. doi: 10.1186/1471-2105-7-S5-S16.
- Wuthrich B, Scheitlin T, Ballmer-Weber B. Isolated allergy to rice. *Allergy.* 2002;57:263–264.
- Xia JF, Wang SL, Lei YK. Computational methods for the prediction of protein-protein interactions. *Protein Pept Lett.* 2010;17:1069–1078. doi: 10.2174/092986610791760405.
- Xiao F, Tang X, Zhou JM. Expression of 35S:: Pto globally activates defense-related genes in tomato plants. *Plant Physiol.* 2001;126:1637–1645. doi: 10.1104/pp.126.4.1637.
- Xie, L., N. Chen, B. Duan, Z. Zhu and X. Liao. 2008. Impact of proteins and cooking properties of waxy and non-waxy rice. *J. Cereal Sci.*, 47: 372-379.
- Xie, Y.R., and N. Hettiarachchy, Xanthan Gum Effects on Solubility and Emulsification Properties. *J. Food Sci.* 62:1101–1104(1997).
- Xu, H., X. Wang and G. Yu, 2006. Optimization of technology for extracting rice bran protein with different enzymes. *Nongye Gongcheng Xuebao/Transac. Chinese Soc. Agric. Eng.*, 22: 157-160.
- Xu C., Li C.Y.T., Kong A.N.T. Induction of phase I, II and III drug metabolism/transport by xenobiotics. *Arch. Pharm. Res.* 2005;28:249–268.
- Xu J., Ge X., Dolan M.C. Towards high-yield production of pharmaceutical proteins with plant cell suspension cultures. *Biotechnol. Adv.* 2011;29:278–299.
- Xu Y, Hanson MR. Programmed cell death during pollination-induced petal senescence in petunia. *Plant Physiol.* 2000;122:1323–1334. doi: 10.1104/pp.122.4.1323.
- Y. Pomeranz, 1992, Effect of drying on rice quality, *Encyclopedia of Food Science and Technology* 1,35.
- Yamada, T., Kato, T., Tamaki, S., Teranishi, K. & Hisamatsu, M. (1998). Introduction of fatty acids to starch granules by ultra-high-pressure treatment. *Starch/Starke*, 11–12, 484–486.

Yamada C, Izumi H, Hirano J, Mizukuchi A, Kise M, Matsuda T, et al. Degradation of soluble proteins including some allergens in brown rice grains by endogenous proteolytic activity during germination and heat-processing. *Biosci Biotechnol Biochem*. 2005;69:1877–1883.

Yamada C, Yamashita Y, Seki R, Izumi H, Matsuda T, Kato Y. Digestion and gastrointestinal absorption of the 14-16-kDa rice allergens. *Biosci Biotechnol Biochem*. 2006;70:1890–1897.

Yang L, Chen JH, Zhang H, Qiu W, Liu QH, Peng X, Li YN, Yang HK: Alkali treatment affects in vitro digestibility and bile acid binding activity of rice protein due to varying its ratio of arginine to lysine. *Food Chem* 2012, 132:925-930.

Yang L, Chen JH, Xu T, Qiu W, Zhang Y, Zhang LW, Xu FP, Liu HB. Rice protein extracted by different methods affects cholesterol metabolism in rats due to its lower digestibility. *Int J Mol Sci*. 2011;12:7594–7608. doi: 10.3390/ijms12117594.

Yang L, Kumagai T, Kawamura H, Watanabe T, Kubota M, Fujimura S, Watanabe R, Kadowaki M: Effects of rice proteins from two cultivars, Koshihikar and Shuny, on cholesterol and triglyceride metabolism in growing and adult rats. *Biosci Biotechnol Biochem* 2007, 71:694-703.

Yang L, Kadowaki M: Effects of rice proteins from two cultivars, Koshihikar and Shuny, on hepatic cholesterol secretion by isolated perfused livers of rats fed cholesterol-enriched diets. *Ann Nutr Metab* 2009, 54:283-290.

Yang L, Kadowaki M: Addition of methionine to rice protein affects hepatic cholesterol output inducing hypocholesterolemia in rats fed cholesterol-free diets. *J Med Food* 2011, 14:445-453.

Yang MY, Peng CH, Chan KC, Yang YS, Huang CN, Wang CJ: The hypolipidemic effect of Hibiscus sabdariff polyphenols via inhibiting lipogenesis and promoting hepatic lipid clearance. *J Agric Food Chem* 2010, 58:850-859.

Yang L, Chen JH, Zhang H, Qiu W, Liu QH, Peng X, Li YN, Yang HK. Alkali treatment affects in vitro digestibility and bile acid binding activity of rice protein due to varying its ratio of arginine to lysine. *Food Chem*. 2012;132:925–930. doi: 10.1016/j.foodchem.2011.11.068.

Yang L., Kumagai T., Kawamura H., Watanabe T., Kubota M., Fujimura S., Watanabe R., Kadowaki M. Effects of rice proteins from two cultivars, Koshihikari and Shunyo, on cholesterol and triglyceride metabolism in growing and adult rats. *Biosci. Biotechnol. Biochem*. 2007;71:694–703.

Yang L., Kadowaki M. Effects of rice proteins from two cultivars, Koshihikari and Shunyo, on hepatic cholesterol secretion by isolated perfused livers of rats fed cholesterol-enriched diets. *Ann. Nutr. Metab*. 2009;54:283–290.

Yano A., Takekoshi M., Morita E., Imai S., Nisizawa T., Hanada N. Production of Fab fragment corresponding to surface protein antigen of Streptococcus mutans serotype c-derived peptide by Escherichia coli and cultured tobacco cells. *J. Biosci. Bioeng*. 2006;101:251–256.

Yasumatsu K; Moritaka, S. and karinum (T. (1964) effect of changes during storages in lipid composition of rice on its amylograph *Agric. B. O. chem*. Vol. 28 pg. 265-272.

- Yeh, A.I. & Li, J.Y. (1996a). Kinetics of phase transition of native, cross-linked, and hydroxypropylated rice starches. *Starch/Starke*, 48, 17–21.
- Yeh, A.I. & Li, J.Y. (1996b). A continuous measurement of swelling of rice starch during heating. *Journal of Cereal Science*, 23, 277–283.
- Yoshino, G., T. Kazumj, M. Amano, M. Tateiwa, T. Yamasakim, S. Takashima, M. Iwai, H. Hatanaka, and S. Baba. 1989. Effects of gamma-oryzanol and probucol on hyperlipidemia. *Current Therap. Res.* 45:975-982.
- Youens-Clark K, Buckler E, Casstevens T, Chen C, Declerck G, Derwent P, Dharmawardhana P, Jaiswal P, Kersey P, Karthikeyan AS. et al. Gramene database in 2010: updates and extensions. *Nucleic Acids Res.* 2010;39:D1085–1094.
- Yu XM, Yu XD, Qu ZP, Huang XJ, Guo J, Han QM, Zhao J, Huang LL, Kang ZS. Cloning of a putative hypersensitive induced reaction gene from wheat infected by stripe rust fungus. *Gene.* 2008;407:193–198. doi: 10.1016/j.gene.2007.10.010.
- Yu HX, Liu QQ, Wang L, Zhao ZP, Xu L, Huang BL, Gong ZY, Tang SZ, Gu NH. Breeding of selectable marker-free transgenic rice lines containing AP1 gene with enhanced disease resistance. *Sci Agric Sin.* 2006;5:805–811.
- Yu H, Braun P, Yildirim MA, Lemmens I, Venkatesan K, Sahalie J, Hirozane-Kishikawa T, Gebreab F, Li N, Simonis N. et al. High-quality binary protein interaction map of the yeast interactome network. *Science.* 2008;322:104–110. doi: 10.1126/science.1158684.
- Yum HY, Lee KE, Choi SY, Yang HS, Sohn MH, Lee SI, et al. Wild rice, hypoallergenic rice, and GMO rice-immuologic comparison. *Pediatr Allergy Respir Dis.* 2005;15:117–125.
- Zanzoni A, Montecchi-Palazzi L, Quondam M, Ausiello G, Helmer-Citterich M, Cesareni G. MINT: a Molecular INTERaction database. *FEBS Lett.* 2002;513:135–140. doi: 10.1016/S0014-5793(01)03293-8.
- Zhai, C.K., Lu, C.M., Zhang, X.Q., Sun, G.J. & Lorenz, K.J. (2001). Comparative study on nutritional value of Chinese and North American wild rice. *Journal of Food Composition and Analysis*, 14, 371–382.
- Zhang G., Malik V.S., Pan A., Kumar S., Holmes M.D., Spiegelman D., Lin X., Hu F.B. Substituting brown rice for white rice to lower diabetes risk: A focus-group study in Chinese adults. *J. Am. Diet. Assoc.* 2010;110:1216–1221.
- Zhang H, Bartley GE, Mitchell CR, Zhang H, Yokoyama W: Lower weight gain and hepatic lipid content in hamsters fed high fat diets supplemented with white rice protein, brown rice protein, soy protein and their hydrolysates. *J Agric Food Chem* 2011, 59:10927-10933.
- Zhang Q, Shi AN, Yang WC, Wang CL. Breeding of three near-isogenic Japonica rice lines with different major genes for resistance to bacterial blight. *Acta Agronom Sin.* 1996;22:135–141.

Zhang W., Hu D., Xi Y., Zhang M., Duan G. Spread of HIV in one village in central china with a high prevalence rate of blood-borne aids. *Int. J. Infect. Dis.* 2006;10:475–480.

Zhang X., Yuan Z., Duan Q., Zhu H., Yu H., Wang Q. Mucosal immunity in mice induced by orally administered transgenic rice. *Vaccine.* 2009;27:1596–1600.

Zhang Y, Gao P, Yuan JS. Plant protein-protein interaction network and interactome. *Curr Genomics.* 2010;11:40–46. doi: 10.2174/138920210790218016.

Zhao Y., Qian Q., Wang H., Huang D. Hereditary behavior of bar gene cassette is complex in rice mediated by particle bombardment. *J. Genet. Genomics.* 2007;34:824–835.

Zhong W, Sternberg PW. Genome-wide prediction of *C. elegans* genetic interactions. *Science.* 2006;311:1481–1484. doi: 10.1126/science.1123287.

Zhou J, Loh Y, Bressan R, Martin G. The tomato gene *Pti1* encodes a serine/threonine kinase that is phosphorylated by *Pto* and is involved in the hypersensitive response. *Cell.* 1995;83:925–935. doi: 10.1016/0092-8674(95)90208-2.

Zhou L, Cheung MY, Zhang Q, Lei CL, Zhang SH, Sun SSM, Lam HM. A novel simple extracellular leucine-rich repeat (eLRR) domain protein from rice (*OsLRR1*) enters the endosomal pathway and interacts with the hypersensitive-induced reaction protein 1 (*OsHIR1*) *Plant Cell Environ.* 2009;32:1804–1820. doi: 10.1111/j.1365-3040.2009.02039.x.

Zhu L, Fu YP, Liu WZ, Hu GC, Si HM, Tang KX, Sun ZX. Rapidly obtaining the marker-free transgenic rice with three target genes by co-transformation an anther culture. *Chin J Rice Sci.* 2007;14:239–247. doi: 10.1016/S1672-6308(08)60001-3. World Health Organization. Diabetes fact sheet. [accessed on 2 June 20] Zobel, H.F. (1988). Molecules to granules: a comprehensive starch review. *Starch/Stärke*, 40, 44–50.12]. Available online: <http://www.who.int/mediacentre/factsheets/fs312/en/index.html>.

Zobel, H.F. (1988). Molecules to granules: a comprehensive starch review. *Starch/Stärke*, 40, 44–50.